

# The warm ionized gas in CALIFA early-type galaxies

## 2D emission-line patterns and kinematics for 32 galaxies

J.M. Gomes<sup>1</sup>, P. Papaderos<sup>1</sup>, C. Kehrig<sup>2</sup>, J. M. Vílchez<sup>2</sup>, M. D. Lehnert<sup>3</sup>, S. F. Sánchez<sup>2,4</sup>, B. Ziegler<sup>5</sup>, I. Breda<sup>1</sup>, S.N. dos Reis<sup>1</sup>, J. Iglesias-Páramo<sup>2,6</sup>, J. Bland-Hawthorn<sup>7</sup>, L. Galbany<sup>8,9</sup>, D. J. Bomans<sup>10,11</sup>, F. F. Rosales-Ortega<sup>12</sup>, R. Cid Fernandes<sup>13</sup>, C. J. Walcher<sup>14</sup>, J. Falcón-Barroso<sup>15,16</sup>, R. García-Benito<sup>2</sup>, I. Márquez<sup>2</sup>, A. del Olmo<sup>2</sup>, J. Masegosa<sup>2</sup>, M. Mollá<sup>17</sup>, R. A. Marino<sup>18,19</sup>, R. M. González Delgado<sup>2</sup>, Á. R. López-Sánchez<sup>20,21</sup>, and the CALIFA collaboration

(Affiliations can be found after the references)

Received ?? ; Accepted ??

### ABSTRACT

**Context.** The morphological, spectroscopic and kinematical properties of the warm interstellar medium (*wim*) in early-type galaxies (ETGs) hold key observational constraints to nuclear activity and the buildup history of these massive, quiescent systems. High-quality integral field spectroscopy (IFS) data with a wide spectral and spatial coverage, such as those from the CALIFA survey, offer an unprecedented opportunity for advancing our understanding of the *wim* in ETGs.

**Aims.** This article centers on a 2D investigation of the *wim* component in 32 nearby ( $\leq 150$  Mpc) ETGs from CALIFA, complementing a previous 1D analysis of the same sample (Papaderos et al. 2013; P13).

**Methods.** The analysis presented here includes  $H\alpha$  intensity and equivalent width (EW) maps and radial profiles, diagnostic emission-line ratios, besides ionized-gas and stellar kinematics. It is supplemented by  $\tau$ -ratio maps as an efficient means to quantify the role of photoionization by the post-AGB stellar component, as compared to alternative mechanisms (e.g., AGN, low-level star formation).

**Results.** Confirming and strengthening our conclusions in P13, we find that ETGs span a broad continuous sequence in the properties of their *wim*, exemplified by two characteristic classes. The first one (type i) comprises systems with a nearly constant  $EW(H\alpha)$  in their extranuclear component, in quantitative agreement with (even though, no proof for) the hypothesis of photoionization by the post-AGB stellar component being the main driver of extended *wim* emission. The second class (type ii) stands for virtually *wim*-evacuated ETGs with a very low ( $\leq 0.5$  Å), outwardly increasing  $EW(H\alpha)$ . These two classes appear indistinguishable from one another by their LINER-specific emission-line ratios in their extranuclear component. Here we extend the tentative classification we proposed previously by the type i+, which is assigned to a subset of type i ETGs exhibiting ongoing low-level star-forming activity in their periphery. This finding along with faint traces of localized star formation in the extranuclear component of several of our sample galaxies points to a non-negligible contribution by OB stars to the global ionizing photon budget in ETGs. Additionally, our data further highlight the diversity of ETGs with respect to their gaseous and stellar kinematics. Whereas, in one half of our sample, gas and stars show similar (yet not necessarily identical) velocity patterns, both dominated by rotation along the major galaxy axis, our analysis also documents several cases of kinematical decoupling between gas and stars, or rotation along the minor galaxy axis. We point out that the generally very low ( $\leq 1$  Å)  $EW(H\alpha)$  of ETGs make necessary a careful quantitative assessment of potential observational and analysis biases in studies of their *wim*. Since, with standard emission-line fitting tools, Balmer emission lines become progressively difficult to detect below an  $EW(H\alpha) \sim 3$  Å, our current understanding of the presence, and 2D emission patterns and kinematics of the diffuse *wim* ETGs may be severely incomplete. We demonstrate that, at the typical emission-line detection threshold of  $\sim 2$  Å in previous studies, most of the extranuclear *wim* emission in an ETG may evade detection, which could in turn prompt its classification as an entirely gas-devoid system.

**Conclusions.** This study adds further observational evidence for a considerable heterogeneity among ETGs with regard to the physical properties and 2D kinematics of their extended *wim* component, and underscores the importance of IFS studies of these systems over their entire optical extent.

**Key words.** galaxies: elliptical and lenticular, cD - galaxies: nuclei - galaxies: ISM - galaxies: kinematics and dynamics - galaxies: star formation

### 1. Introduction

The featureless appearance of early-type galaxies (ETGs) has for decades sustained the view that these systems are largely ‘simple’, both in terms of their mass assembly history and kinematics, formed early on, and having undergone little evolution over the past several Gyr (e.g., Mathews & Baker, 1971; Bregman, 1978; White & Chevalier, 1983). However, our understanding of the buildup of these systems has undergone a substantial revision over the last years, as ground-based and satellite multi-wavelength data have gradually revealed a great deal

of structural and kinematical diversity in present-day ETGs (e.g., Bender et al., 1989; Kormendy et al., 2009).

For instance, careful photometric studies have revealed in many of these systems a rich variety of low-surface brightness non-axis-symmetric features, such as ripples and shells (e.g., Schweizer & Seitzer, 1988; Struck, 1999, and references therein), pointing to a tumultuous assembly history, with multiple minor merger episodes, some of which may have provided the gas fuel for rejuvenating these ‘old and dead’ galaxies with low-level star formation (SF) (see, e.g., Trager et al., 2000; Rampazzo et al., 2007; Kaviraj et al., 2008; Clemens et al., 2009; Thomas et al., 2010, and references therein). Indeed, the UV-to-

Send offprint requests to: Jean Michel Gomes; jean@astro.up.pt

optical colors of several ETGs are consistent with a prolonged phase of low-level SF, in which 1% to 3% of the stellar mass  $M_*$  could have formed in the past one Gyr (Kaviraj et al., 2007).

With the advent of integral field spectroscopy (IFS), spatially resolved analyses have further highlighted the diversity and complexity of ETGs. For instance, following the initial study of NGC 3377 by Bacon et al. (2001), the SAURON project has given a decisive contribution to the systematization of the kinematical properties of ETGs, among others, by uncovering kinematically decoupled or even counter-rotating cores in several of these systems (e.g., Sarzi et al., 2006). Further insights into the connection between kinematics and other ETG properties are being added with the ATLAS<sup>3D</sup> IFS survey (Cappellari et al., 2011), supplementing previous work with information on, e.g., the relation between the specific angular momentum and the shape of surface brightness profiles (e.g., Krajnovic et al., 2013), or that between the degree of rotational support and X-ray luminosity (Sarzi et al., 2012).

Another key insight has been the presence of faint nebular emission in the majority of ETGs studied with longslit and single-aperture SDSS spectroscopy (e.g., Phillips et al., 1986; Demoulin-Ulrich et al., 1984; Kim, 1989; Trinchieri & di Serego Alighieri, 1991; Annibali et al., 2010; Yan & Blanton, 2012) or narrow-band imaging (Finkelman et al., 2010; Kulkarni et al., 2014). In particular, SAURON, has detected extended nebular emission in  $\sim 75\%$  of the studied ETGs (48 in total), strengthening previous evidence for the presence of an ubiquitous warm ( $T \sim 10^4$  K) interstellar medium (*wim*) component in these systems, and adding new information on its kinematics. For example, Sarzi et al. (2006, see also Sarzi et al. 2007) reported the detection of extended Balmer H $\beta$  emission in ETGs down to an equivalent width (EW) of  $\approx 0.1$  Å, and demonstrated a puzzling variety of morphologies and kinematics, with striking cases of kinematical decoupling between gas and stellar motions.

Using IFS data from the *Calar Alto Legacy Integral Field Area* survey (CALIFA, Sánchez et al., 2012), the presence of an extended *wim* component in ETGs was first demonstrated by Kehrig et al. (2012, hereafter K12), and subsequently by Papaderos et al. (2013) and Singh et al. (2013). CALIFA is the first IFS survey simultaneously permitting full coverage of the optical spectral range and spatial extent of a representative sample of Hubble-type galaxies in the local universe (see Walcher et al., 2014, for description of the sample selection). The wide spectral and spatial coverage of CALIFA is a key advantage over SAURON and ATLAS<sup>3D</sup>, which target a narrow spectral interval encompassing the H $\beta$  and [O III]5007 lines over a field of view (FoV) of  $33'' \times 41''$  ( $\sim 1/3$  of that of CALIFA). As recently pointed out by Arnold et al. (2014) and Gomes et al. (2015b), a full mapping of the optical galaxy size is essential for an unbiased determination of the properties of gas and stars in ETGs.

Spatially resolved IFS studies of emission-line EWs and line-flux ratios of the *wim* also yield key constraints on the energy balance and gas excitation mechanisms in ETGs – an issue being debated for at least the last three decades. Most ETG nuclei show faint nebular emission with an H $\alpha$  equivalent width of typically  $\text{EW}(\text{H}\alpha) \leq 10$  Å (see, e.g., Annibali et al., 2010) and are classified as *low-ionization nuclear emission-line regions*<sup>1</sup> (LINERs; Heckman, 1980) on the basis of optical diagnostic

line ratios after Baldwin et al. (1981, referred to in the following as BPT ratios) and (Veilleux & Osterbrock, 1987, hereafter VO). One widely favored mechanism for the *wim* excitation in ETG nuclei involves ‘weak’, *low-luminosity active galactic nuclei* (LLAGN) (cf, e.g., Ho, 1999), powered by radiatively inefficient (sub-Eddington) accretion onto a central super-massive black hole (SMBH), a process that is thought to be governed by the absence of the broad-line region and its surrounding torus, as in classical Active Galactic Nuclei (AGN) (Ho, 2008, and references therein).

Other hypotheses involve fast shocks (e.g., Dopita & Sutherland, 1995; Allen et al., 2008), photoionization by ongoing low-level SF (e.g., Schawinski et al., 2007; Shapiro et al., 2010) and hot evolved ( $\geq 10^8$  yr) post-asymptotic giant branch (pAGB) stars (e.g., Trinchieri & di Serego Alighieri, 1991; Binette et al., 1994; Macchetto et al., 1996; Stasińska et al., 2008; Cid Fernandes et al., 2010, 2011; Sarzi et al., 2010; Yan & Blanton, 2012). The latter mechanism has attracted a great deal of interest in the last years and was even proposed to be the sole source of gas excitation in ETG/LINERs. Turning into this subject, following earlier work by di Serego Alighieri et al. (1990) and Trinchieri & di Serego Alighieri (1991), Binette et al. (1994) have first studied quantitatively with evolutionary synthesis models the ionizing output from white dwarfs and hot pAGB stars<sup>2</sup>, and demonstrated that these sources are capable of producing the ionizing radiation for simultaneously explaining the low nuclear EWs and LINER characteristics of ETG nuclei. Several observational studies have added support to the pAGB photoionization hypothesis. For example, Sarzi et al. (2010), using SAURON IFS data, conclude that pAGB stars, instead of fast shocks, are the main source of ionizing photons in ETGs. This view was reinforced, at least as far extranuclear *wim* emission is concerned, by our study in K12, and, more recently, by a combined study of Sloan Digital Sky Survey (SDSS; York et al., 2000) and Palomar survey (Ho et al., 1995) spectra by Yan & Blanton (2012). Likewise, Eracleous et al. (2010) point out that pAGB stars provide more ionizing photons than AGN in more than half of the LINERs and can account for the observed line emission in 1/3 of these systems. On the other hand, arguments against the predominance of pAGB photoionization in ETGs/LINERs were discussed in other studies. For instance, Annibali et al. (2010) conclude from the analysis of long-slit spectra for 65 ETGs that in only  $\sim 1/5$  of their sample can the nuclear LINER emission be attributed to pAGB photoionization alone, and excitation by a low-accretion rate AGN or fast shocks is likely necessary for the majority of ETG nuclei. The pAGB component as the dominant source of photoionization in ETG nuclei has also been disputed by, e.g., Ho (2008), on the basis of various arguments, one of them being that line emission in these systems tends to be very centrally concentrated.

Following the seminal theoretical work by Binette et al. (1994), the pAGB photoionization hypothesis has been recently further investigated in e.g., Sodr  & Stasińska (1999); Stasińska et al. (2008); Cid Fernandes et al. (2010, 2011); Stasińska et al. (2015) and confronted with SDSS spectroscopic data. As Stasińska et al. (2008) argue, a significant fraction of ETGs are in fact ‘retired’ galaxies, i.e. systems that stopped forming stars, and whose ionizing field is powered solely by pAGB sources. According to the classification by Cid Fernandes et al. (2011), this large population of retired ETG/LINERs is characterized by

<sup>1</sup> Since it is now recognized that LINER-specific emission-line ratios are not confined to galaxy nuclei (e.g., Sharp & Bland-Hawthorn, 2010; Kehrig et al., 2012; Papaderos et al., 2013; Singh et al., 2013), a more adequate acronym would be *low-ionization emission-line regions* (LIERS).

<sup>2</sup> For the sake of brevity we will refer to in the following as ‘pAGB stars’ to all ionizing point sources arising at and beyond the post-AGB phase.

a  $[\text{N II}]/\text{H}\alpha$  ratio of  $\geq 1$  (similar to classical AGN) yet faint nebular emission ( $0.5 \geq \text{EW}(\text{H}\alpha) [\text{\AA}] \lesssim 3.0$ ). Quite importantly, the generally satisfactory agreement between pAGB photoionization models and single-aperture SDSS data has called into question the necessity of any additional energy source (SF, AGN) in ETGs, and led even to the rejection of ‘weak’ (low-accretion rate, low-luminosity) AGN in these systems: As emphatically pointed out in [Cid Fernandes et al. \(2011\)](#), retired galaxies (i.e. low- $\text{EW}(\text{H}\alpha)$  ETG/LINERs) are actually fake AGN, *erroneously counted as AGN, and leading to the illusion of a Seyfert-LINER dichotomy in the AGN population*.

In the framework of the CALIFA collaboration, two recent studies (P13, [Singh et al., 2013](#)) have confirmed and strengthened the conclusion by K12 that LINER-like nebular emission in ETGs typically extends several kpc away from the galaxy nuclei. They also lent further support, on the basis of distinctively different arguments, to the idea that pAGB photoionization can be an important ingredient of the *wim* excitation on galactic scales. More specifically, P13 studied the radial distribution of the  $\text{EW}(\text{H}\alpha)$  over nearly the entire optical extent of 32 ETGs and demonstrated that this quantity shows a nearly constant value of  $\sim 1 \text{ \AA}$  in the extranuclear component of  $\sim 40\%$  of their sample (type i ETGs in their classification), in quantitative agreement with the diffuse floor of ionization expected from the pAGB background. A second test made involves the  $\tau$  ratio (K12, see also [Binette et al., 1994](#); [Cid Fernandes et al., 2011](#), for similar definitions), defined as the inverse ratio of the observed  $\text{H}\alpha$  luminosity to that predicted from pAGB photoionization when assuming standard conditions in the gas and case B recombination. The nearly constant  $\tau \approx 1$  determined in their extranuclear component of type i ETGs implies that the Lyman continuum photon rate from pAGB stars is energetically capable of sustaining the observed low-level *wim* emission.

On the other hand, P13 have shown that ETGs in their majority display more complex 2D gas emissivity patterns than what is expected from in situ (case B) recombination of the  $\text{Ly}_c$  output from the pAGB component. Specifically, in  $\sim 60\%$  of the ETGs in their sample (type ii in their classification) the  $\tau$  ratio was determined to be between  $\geq 2$  and up to  $\gtrsim 50$ , implying that the bulk of  $\text{Ly}_c$  photons produced by pAGB stars escapes without being locally reprocessed into nebular emission. They interpreted this and the positive radial  $\text{EW}(\text{H}\alpha)$  gradients of type ii ETGs as the result of the interplay between various physical parameters determining the 3D characteristics of the warm and hot gas component in ETGs, such as, e.g., porosity, filling factor and ionization parameter. The fact that type i and type ii ETGs are almost indistinguishable from one another by their LINER characteristics, despite a difference of  $\geq 1$  dex in their  $\tau$  (hence, also their 3D gas structure and  $\text{Ly}_c$  photon escape fraction) has led P13 to conjecture that, in the 3D geometry of ETGs, classical BPT diagnostics become degenerate over the area of the diagrams which cover galaxies with LINER-like ratios. Additionally, as first pointed out in P13, a far-reaching implication of the large  $\text{Ly}_c$  escape fraction in type ii ETGs is that these systems could host significant accretion-powered nuclear activity that evades detection through optical spectroscopy. This is because, in the presence of extensive  $\text{Ly}_c$  escape, the luminosities of nebular emission lines commonly used as diagnostics of accretion-powered nuclear activity and/or to which SMBHs mass determinations are tied, are reduced by at least one order of magnitude. For the same reason, a low nuclear  $\text{EW}(\text{H}\alpha)$  ( $0.5 \dots 3 \text{ \AA}$ ) is not per se compelling evidence for the *wim* emission in ETGs being solely powered by pAGB stars, even though their ionizing photon output is able to

sustain their diffuse  $\text{H}\alpha$  emission. As we recently pointed out in [Gomes et al. \(2015a\)](#), it is imaginable that the combined effect of different gas excitation sources (SF, pAGB, AGN and shocks) with geometry-dependent  $\text{Ly}_c$  leakage can lead to a nearly constant  $\text{EW}(\text{H}\alpha) \sim 1 \text{ \AA}$ , mimicking a predominance of pAGB photoionization all over the extranuclear component of type i ETGs.

Large-FoV, wide spectral coverage IFS data, as those from CALIFA, obviously offer a tremendous potential for further elucidating the nature and excitation mechanisms of the *wim* in ETGs. In this article, we discuss in more detail our methodology in P13 and extend our analysis by a concise summary of the 2D properties of the nebular and stellar component in our ETG sample, as derived with our IFS data processing pipeline PORTO3D. This paper is organized as follows: Section 2 provides a brief outline of the CALIFA survey and basic information on the studied ETG sample. In Sect. 3 we briefly summarize our analysis methodology and in Sect. 4 we describe the output from PORTO3D and auxiliary codes that is considered in the subsequent analysis. In Sect. 5 we provide a comparative discussion of the main properties of type i and type ii ETGs (Sects. 5.2&5.3) laying emphasis on the morphology and EW distribution of their nebular component. The incidence of accretion-powered nuclear activity and the kinematical properties of gas and stars in our sample ETGs are cursorily discussed in Sects. 5.3.1 and Sect. 5.4, respectively. Finally, in Sect. 5.5 we briefly comment on how observational bias stemming from an EW detection threshold for faint nebular emission could affect spectroscopic classifications of ETGs. The main results and conclusions from this study are summarized in Sect. 6.

The Appendix provides theoretical predictions from the literature and our own evolutionary synthesis code on the time evolution of the  $\text{EW}(\text{H}\alpha)$  for a pAGB-dominated stellar component (Sect. A). Additionally, in Sect. B we discuss the main components of PORTO3D, and the methodology employed for the determination of emission-line fluxes and their uncertainties, as well as the derivation of the radial distribution of various quantities of interest. This article is supplemented by the main output from PORTO3D for our sample ETGs (Sect. C) and a comparison sample of composite/star-forming galaxies from CALIFA (Sect. D), which is meant to illustrate the variation of the  $\text{EW}(\text{H}\alpha)$ ,  $\tau$  and diagnostic line ratios from early-type towards late-type galaxies.

## 2. Data sample

The CALIFA survey ([Sánchez et al., 2012](#), see also Sánchez 2014a for a review) aims at a systematic study of a representative sample of the Hubble-type galaxy population in the local Universe ( $z \leq 0.03$ ). The CALIFA sample (600 galaxies, [Walcher et al., 2014](#)) has been selected such as to ensure optimal statistics for the different Hubble types, and to take maximum advantage of the large hexagonal FoV ( $74'' \times 62''$ , sampled by 331 fibers 36 of which are dedicated to sky background subtraction) of the PMAS/PPak spectrograph ([Roth et al., 2005](#); [Kelz et al., 2006](#)) mounted at the 3.5m telescope at the Calar Alto observatory. For each sample galaxy, the CALIFA IFS data combine 2 to 3 individual dithered exposures in two different instrument setups: the low-spectral resolution setup (V500;  $R \sim 850$ ) and the medium-resolution setup (V1200;  $R \sim 1700$ ) with a spectral coverage between  $3750 - 7500 \text{ \AA}$  and  $3700 - 4200 \text{ \AA}$ , respectively. After reduction with the CALIFA IFS data processing pipeline (version 1.3c for the data set used here; see [Husemann et al., 2013](#), for details), the data are provided to



the CALIFA collaboration interpolated within a  $78'' \times 72''$  grid, flux-calibrated at a precision better than  $\sim 15\%$  and corrected for foreground Galactic extinction. Additionally, the CALIFA data processing pipeline provides from its version v1.3c onward error statistics for each spaxel and a detailed quality control (e.g., spectral masks with spurious features, such as local residuals in the sky- or cosmic-ray removal). Regular public data releases of fully reduced CALIFA IFS data, the latest one (DR2; 200 galaxies – [García-Benito et al., 2015](#)) in fall 2014, underscore the legacy potential of the CALIFA survey and its general value to the astronomical community.

Following its first science publication in K12, the CALIFA collaboration has completed several studies dedicated to, e.g., aperture effects ([Iglesias-Páramo et al., 2013](#)) and the spatial resolution of IFS data ([Mast et al., 2013](#)), stellar age gradients in galaxies ([Pérez et al., 2013](#); [González Delgado et al., 2014a](#); [Sánchez-Blázquez et al., 2014](#)) and the uncertainties in their determination from spectral fitting ([Cid Fernandes et al., 2014](#)), the mass-metallicity relation ([Sánchez et al., 2013](#); [González Delgado et al., 2014b](#)), the presence of universal metallicity gradients in galaxies ([Sánchez, 2014a](#); [Sánchez et al., 2014b](#)), the spatial distribution and excitation mechanisms of nebular emission in ETGs and other LINER galaxies (P13, [Singh et al., 2013](#)), and devised a new semi-empirical metallicity calibration based on the largest hitherto compiled sample of HII regions in late-type galaxies ([Marino et al., 2013](#)). Several parallel ongoing studies within the collaboration explore various properties of present-day Hubble-type galaxies, such as their stellar and gas kinematics ([García-Lorenzo et al., 2015](#); [Barrera-Ballesteros et al., 2014](#), Falcón-Barroso et al., in prep.).

This study is based on V500 IFS datacubes for 20 E and 12 S0/SA0 galaxies in the local volume ( $<150$  Mpc) and comprises essentially all ETGs observed by CALIFA at the onset of this project (mid 2012) with our pilot study in K12. The low-resolution (V500) CALIFA IFS data have the advantage of covering the entire optical spectral range, thereby permitting flux determinations of several emission lines and their ratios, and spectral fitting of single-spaxel spectra down to a surface brightness level  $\mu \simeq 23.6$  g mag/arcsec<sup>2</sup> (cf, e.g., K12), allowing for the study of the extranuclear component of ETGs out to typically  $\gtrsim 2$   $r$  band Petrosian<sub>50</sub> radii  $r_p$ , taken from SDSS.

In P13 we studied these 32 ETGs (Table 1) only with respect to the radial distribution of their EW(H $\alpha$ ),  $\tau$  and BPT ratios, without discussing the 2D properties of their nebular and stellar components, which is the goal of the present article.

### 3. Outline of the analysis methodology

The CALIFA IFS data were processed spaxel-by-spaxel with PORTO3D, a pipeline developed by us with the purpose of automated spectral fitting and post-processing of flux-calibrated IFS data cubes. This pipeline (see Fig. B.1 for a schematic overview) combines a suite of modules written in the MIDAS<sup>3</sup> script language and in GNU Fortran 2008, with peripheral modules using CFITSIO and PGplot routines. The spectral fitting module of PORTO3D invokes the publicly available (v.4) version of the population synthesis code STARLIGHT ([Cid Fernandes et al., 2005](#)). PORTO3D enabled the first science publication of the CALIFA collaboration in K12 and since then has undergone several upgrades. Its current version (v.2), used in P13 and this

study, incorporates several additions, in particular, it provides estimates of uncertainties in emission-line fluxes and EWs.

The pipeline consists of three main modules: The first one (m.1) performs a data quality assessment and initial statistics, and extracts from the flux-calibrated CALIFA IFS data cubes individual spectra, which in turn are transformed into a format suitable for fitting with STARLIGHT. Module m.2 is intended to the extraction of emission lines, after subtraction of the best-fitting stellar fit, and the determination of emission-line fluxes and kinematics. It also computes various secondary quantities from the STARLIGHT models (e.g., luminosity- and mass-weighted stellar age and metallicity, the luminosity fraction of stars younger than 5 Gyr, and others). The third module, m.3, determines the Balmer line luminosities implied by the  $L_{Y_c}$  output from the best-fitting population vector. The predicted H $\alpha$  luminosity is in turn compared with the observed one in order to assess via the  $\tau$  ratio (K12,P13) the consistency of the results with the pAGB photoionization hypothesis.

Each ETG has been processed with PORTO3D twice, based on two different libraries of Simple Stellar Population (SSP) models that were used for spectral modeling with STARLIGHT. The emission-line maps produced by these two runs were in turn combined by a decision-tree routine, which permits check of the soundness and subsequent error-weighted averaging of values spaxel-by-spaxel. PORTO3D is supplemented by a routine for computation of radial profiles of various quantities of interest (e.g., emission-line intensities and EWs,  $\tau$  and BPT ratios). This add-on module is essentially an adaptation of the isophotal annuli surface photometry technique iv by ([Papaderos et al., 2002](#), hereafter P02).

<sup>3</sup> Munich Image Data Analysis System, provided by the European Southern Observatory (ESO).



**Table 1.** General properties of the sample ETGs.

Name (1)	Morph. (2)	T (3)	$\alpha$ (J2000) (4)	$\delta$ (J2000) (5)	D (6)	$m_r$ (7)	$r_p$ (8)	$f_{H\alpha}$ (9)	$L_{H\alpha}$ (10)	$M_*$ (e/p) (11)	$\langle\tau\rangle$ (12)	SC (13)	KC (14)
UGC 5771	S0/a	i	10:37:19.33	+43:35:15.31	106.4	13.23	7.3	251	34.0	11.560/11.398	0.69	L	r  r
NGC 4003	SB0	i	11:57:59.03	+23:07:29.63	96.6	13.21	9.6	455	50.8	11.378/11.218	0.57	C	r  p
UGC 8234	S0/a	i	13:08:46.50	+62:16:18.10	116.1	12.82	5.2	99	15.9	11.270/11.131	1.09	L	r  n
NGC 5966	E	i	15:35:52.10	+39:46:08.05	69.0	12.52	9.4	206	117.2	11.311/11.155	1.26	L	r  o
UGC 10205	Sa	i	16:06:40.18	+30:05:56.65	97.6	13.16	12.2	561	63.9	11.375/11.221	0.20	C/L	r  p
NGC 6081	S0	i	16:12:56.85	+09:52:01.57	79.6	12.86	8.2	249	18.9	11.493/11.331	1.07	L	r  p
NGC 6146	E?	i	16:25:10.32	+40:53:34.31	127.4	12.49	7.2	180	34.9	11.880/11.720	1.86	L	r  p
UGC 10695	E	i	17:05:05.57	+43:02:35.35	120.1	13.19	10.7	274	47.3	11.493/11.335	0.90	L(S3)	n+o
UGC 10905	S0/a	i	17:34:06.43	+25:20:38.29	114.1	13.03	7.1	217	33.8	11.641/11.487	0.98	L	r  p
NGC 6762	S0/a	i	19:05:37.09	+63:56:02.79	44.9	13.19	5.8	236	56.7	11.678/11.524	0.77	L	r  p
NGC 7025	Sa	i	21:07:47.33	+16:20:09.22	70.7	12.20	9.2	604	36.1	11.655/11.496	1.67	L	r  p
NGC 1167	SA0	i+	03:01:42.33	+35:12:20.21	66.2	12.19	13.6	827	43.3	11.572/11.410	0.87	L(S3)●1	r  p
NGC 1349	S0	i+	03:31:27.51	+04:22:51.24	87.7	13.09	11.7	467	42.9	11.341/11.183	0.89	L	r  p
NGC 3106	S0	i+	10:04:05.25	+31:11:07.65	90.1	12.69	11.0	651	63.0	11.474/11.314	0.60	L	r  p
UGC 29	E	ii	00:04:33.74	+28:18:06.20	118.5	13.44	8.9	33	5.6	11.362/11.203	4.98	L	n+n
NGC 2918	E	ii	09:35:44.04	+31:42:19.67	98.1	12.56	8.3	60	6.9	11.501/10.340	4.87	L	r  n
NGC 3300	SAB(r)0	ii	10:36:38.44	+14:10:15.97	48.0	12.45	10.9	39	1.1	11.012/10.858	19.8	L	r  n
NGC 3615	E	ii	11:18:06.65	+23:23:50.36	98.3	12.50	6.9	35	4.0	11.690/11.523	10.8	L	r  n
NGC 4816	S0	ii	12:56:12.14	+27:44:43.71	102.6	12.91	12.7	35	4.4	11.530/11.367	18.6	L	p  n
NGC 6125	E	ii	16:19:11.53	+57:59:02.89	72.2	12.13	9.1	52	3.2	11.600/11.437	23.4	L	p+n
NGC 6150	E?	ii	16:25:49.96	+40:29:19.41	126.0	13.12	7.6	20	3.8	11.660/11.496	10.9	L●2	r  n
NGC 6173	E	ii	16:29:44.87	+40:48:41.96	126.9	12.33	11.0	45	8.6	12.161/11.999	15.6	L	r  n
UGC 10693	E	ii	17:04:53.01	+41:51:55.76	120.5	12.70	9.3	34	5.9	11.731/11.572	11.7	L	p  n
NGC 6338	S0	ii	17:15:22.97	+57:24:40.28	117.5	12.56	11.3	178	29.4	11.870/11.707	2.21	L	r  p
NGC 6411	E	ii	17:35:32.84	+60:48:48.26	57.6	12.09	11.3	97	3.9	11.363/11.203	8.62	C/L	p  p
NGC 6427	S0	ii	17:43:38.59	+25:29:38.18	51.3	12.71	4.9	68	2.1	11.141/10.985	7.60	L	r  n
NGC 6515	E	ii	17:57:25.19	+50:43:41.24	99.0	12.87	9.2	87	10.2	11.474/11.316	2.46	L/S	n+n
NGC 7194	E	ii	22:03:30.93	+12:38:12.41	110.7	12.87	6.6	30	4.4	11.698/11.536	12.9	L	r  n
NGC 7236	SA0	ii	22:14:44.98	+13:50:47.46	108.2	13.72	3.4	35	5.0	11.565/11.403	8.16	L	r  n
UGC 11958	SA0	ii	22:14:46.88	+13:50:27.13	108.3	13.35	14.2	153	21.4	11.776/11.612	2.67	L●3	n+n
NGC 7436B	E	ii	22:57:57.54	+26:09:00.01	100.6	12.80	10.0	94	11.4	11.867/11.708	6.87	L	r  n
NGC 7550	SA0	ii	23:15:16.07	+18:57:41.22	69.2	12.29	11.6	267	15.3	11.555/11.394	3.02	L	n+r
IC 540	S	–	09:30:10.33	+07:54:09.90	31.9	13.61	8.9	401	4.9	10.152/10.004	0.36	S	r  p
UGC 8778	S?	–	13:52:06.66	+38:04:01.27	52.6	13.56	9.6	590	19.6	10.590/10.439	0.31	C/L	r  r
IC 944	Sa	–	13:51:30.86	+14:05:31.95	104.8	12.89	10.6	530	70	11.556/10.439	0.58	C/L	r  p
IC 1683	S?	–	01:22:38.92	+34:26:13.65	65.4	13.47	9.2	1703	87	10.924/10.772	0.18	H	r  p
NGC 4470	Sa?	–	12:29:37.77	+07:49:27.12	38.4	12.57	13.3	8372	147	10.231/10.108	0.03	H	r  r

★ Col. 1: Galaxy name.

★ Col. 2: Morphological classification adopted from NASA/IPAC Extragalactic Database (NED).

★ Col. 3: Classification according to our study in P13, based on the mean  $\tau$  ratio (Col. 12) for the extranuclear component (type i, i+ and ii).

★ Col. 4: Right ascension.

★ Col. 5: Declination.

★ Col. 6: Distance in Mpc adopted from NED.

★ Col. 7: Apparent  $r$  total magnitude from SDSS DR7.★ Col. 8: Petrosian\_50 radius in arcsec in the SDSS  $r$  band.★ Col. 9: Integral  $H\alpha$  flux in units of  $10^{-16} \text{ erg s}^{-1} \text{ cm}^{-2}$ , derived from CALIFA IFS data.★ Col. 10: Integral  $H\alpha$  luminosity in units of  $10^{39} \text{ erg s}^{-1}$ .★ Col. 11: Logarithm of the ever formed and present stellar mass in  $M_\odot$ , as obtained from spectral modeling with STARLIGHT.★ Col. 12: Mean  $\tau$  ratio (from P13 in the case of ETGs).★ Col. 13: Spectroscopic classification based on the emission-line ratios (panels i&j of Figs. 1, C.1–C.31 and D.1–D.5) of the nuclear ( $R^* \leq 3''$ ) CALIFA spectrum. The notation is as follows: **H**: HII/star-forming, **C**: composite (cf description of panels i&j of Fig. 1), **L**: LINER, **S**: Seyfert. Whenever available, the spectroscopic classification from NED is included within brackets, and the index following the ● symbol points to additional information available at NED on the morphology of the radio continuum: (1) Radio jet, (2) Narrow-angle tail radio source, (3) FRI I source.★ Col. 14: Tentative 3-character kinematical classification: **r**: rotation-dominated kinematics, **p**: perturbed rotational pattern, **o**: gas outflow, and **n**: no distinguishable pattern or pressure-supported stellar kinematics. The first and third character refer, respectively, to the kinematics of stars and gas, and the second one indicates in the case of stellar rotation (r or p) whether the kinematic and photometric major axes are roughly aligned (||) or not (||). The + symbol marks ETGs where stellar rotation is not detected (n) or having a nearly circular-symmetric morphology, thus uncertain photometric major axis.Note that the galaxies NGC 7025, NGC 7236 and NGC 7436B are overlapping over a substantial fraction of their area with extended foreground/background sources that, even after subtraction, are likely introducing uncertainties of  $\sim 10\%$  in the quoted  $H\alpha$  fluxes.

## 4. Results

Figure 1 shows a representative example of the information extracted and used in the discussion of our ETG sample. This figure along with the results obtained for the remaining sample galaxies (Figs. C.1–C.31; Appendix C) illustrate the considerable diversity of local ETGs with respect to morphology, kinematics and physical properties of their warm interstellar medium. The 2D maps (panels a–f & n) cover an area of  $78'' \times 72''$  and are displayed in astronomical orientation (north is up and east to the left).

Panel a: Logarithmic representation of the emission-line free stellar continuum flux (in units of  $10^{-16} \text{ erg s}^{-1} \text{ cm}^{-2}$ ) in the spectral range between 6390 Å and 6490 Å. The cross (in panels a–f and n) marks the maximum intensity of the stellar emission, the morphology of which is further delineated by the overlaid contours at 32, 10, 3, 2.3 and 1.3% of the peak intensity. The linear scale in kpc labeling the  $20''$  horizontal bar to the upper-right has been computed from the adopted distance to the source (Col. 6 in Table 1). Regions being contaminated by probable foreground or background sources, therefore excluded from the analysis, are left blank.

Panel b:  $H\alpha$  flux map in units of  $10^{-16} \text{ erg s}^{-1} \text{ cm}^{-2}$ .

Panel c:  $EW(H\alpha)$  map in Å.

Panel d: Stellar velocity field after correction for systemic velocity, as derived from STARLIGHT fits to individual spaxels. The right-hand side vertical bar indicates the dynamical range of the image in  $\text{km s}^{-1}$ , and the red circle centered on the linear scale bar illustrates the effective angular resolution ( $\text{FWHM} \approx 3''.8$  for data cubes processed with version 1.3c of the CALIFA data reduction pipeline) of our IFS data. The blue lines show the photometric minor axis.

Panel e: Ionized gas velocity field, as obtained by averaging the  $H\alpha$  and  $[\text{NII}]\lambda 6584$  velocity maps.

Panel f: Subdivision of the  $EW(H\alpha)$  map into three intervals, meant to help the eye to distinguish between regions where the observed  $EW(H\alpha)$  is consistent with pure pAGB photoionization ( $0.5\text{--}2.4$  Å; light blue), an additional gas excitation source is needed to account for the observed  $EW(H\alpha)$  ( $>2.4$  Å; orange), and where the  $EW(H\alpha)$  is lower than that predicted from pAGB photoionization models ( $\leq 0.5$  Å; dark blue; cf Sect. A), thus  $Ly_c$  photon escape is likely important. The fraction in % of the spectroscopically studied area where the observed  $EW(H\alpha)$  implies  $Ly_c$  leakage, is consistent with pure pAGB photoionization, or an extra ionization source is required is indicated in the upper-right of this panel. Note that the adopted lower bound of  $0.5$  Å ( $\equiv EW_{\star}^{-}$ ) corresponds to the  $EW(H\alpha)$  expected for an instantaneous burst with an age of  $\sim 1$  Gyr (cf evolutionary synthesis models in Sect. A), i.e. it is rather representative of ETGs having undergone significant recent stellar mass growth. Since the  $EW(H\alpha)$  expected for a typical ETG (age  $\geq 5$  Gyr) is  $\geq 1$  Å, the condition  $EW(H\alpha) \geq 0.5$  Å is rather a generous minimum requirement for the validity of the pAGB photoionization hypothesis, provided that  $Ly_c$  escape is negligible. Panels g&h:  $H\alpha$  intensity, normalized to its peak value, and  $EW(H\alpha)$  as a function of the photometric radius  $R^*$  ( $''$ ). Determinations based on single spaxels (sisp) within the nuclear region, defined as twice the angular resolution of the IFS data (i.e. for  $R^* \leq 3''.8$ ) and in the extranuclear component are shown with red and blue circles, respectively. Open squares (green color) correspond to the average of individual sisp determinations within isophotal annuli (isan; cf Sect. B.4) with vertical bars illustrating the  $\pm 1\sigma$  scatter of data points. The light-shaded area illustrates the  $EW(H\alpha)$  range that can be accounted for by pAGB photoionization for ages between

$\sim 0.1$  Gyr and  $\sim 13$  Gyr for the full range in metallicity covered by SSP models from Bruzual & Charlot (2003, hereafter BC03). This range includes a brief minimum ( $0.1\text{--}0.8$  Å) at the onset of the pAGB phase and an upper range of  $3.4$  Å for a super-solar metallicity of  $2.5 Z_{\odot}$  ( $0.1\text{--}3.4$  Å; cf Fig. A.1 in this paper and Fig. 2 in Cid Fernandes et al. (2011)). However, for reasonable assumptions on the age ( $1\text{--}12$  Gyr) and metallicity ( $\sim Z_{\odot}$ ) of stellar populations in ETGs (Sect. A for details), the range of expected  $EW(H\alpha)$ 's can be narrowed down to within  $EW_{\star}^{-} = 0.5$  Å and  $EW_{\star}^{+} = 2.4$  Å (dark-gray strip) with a typical value around  $1$  Å ( $\equiv \langle EW_{\star} \rangle$ ) in the age interval between  $\sim 6$  Gyr and  $\sim 11$  Gyr.


Panels i&j: sisp determinations for the BPT-VO diagrams on the  $\log([\text{N II}]\lambda 6584/H\alpha)$  vs  $\log([\text{O III}]\lambda 5007/H\beta)$  and  $\log(\Sigma[\text{S II}]\lambda 6717, 6731/H\alpha)$  vs  $\log([\text{O III}]\lambda 5007/H\beta)$  diagnostic emission-line ratios planes. The meaning of symbols is identical to those in panels g–h. The loci on the BPT diagrams that are characteristic of AGN and LINERs, and that corresponding to photoionization by young massive stars in HII regions are indicated, with demarcation lines from Kauffmann et al. (2003, dotted curve), Kewley et al. (2001, solid curve) and Schawinski et al. (2007, dashed line). For the sake of comparison, the grid of thin-gray lines roughly at the middle of each diagram depicts the parameter space that can be accounted for by pure shock excitation, as predicted by Allen et al. (2008) for a magnetic field of  $1 \mu\text{G}$ , and a range of shock velocities between  $100$  and  $1000 \text{ km s}^{-1}$ , for gas densities between  $0.1$  and  $100 \text{ cm}^{-3}$ .

Panels k&l:  $\log([\text{O III}]\lambda 5007/H\beta)$  and  $\log([\text{N II}]\lambda 6584/H\alpha)$  ratio as a function of  $R^*$ , as inferred from sisp and isan determinations. The shaded horizontal area depicts the mean ratios for our sample (cf P13) of  $0.37 \pm 0.13$  for  $\log([\text{O III}]/H\beta)$  and  $0.34 \pm 0.26$  for  $\log([\text{N II}]/H\alpha)$ , with a standard deviation about the mean  $\sigma_N$  of  $0.02$  and  $0.05$ , respectively.

Panel m:  $\tau$  ratio radial distribution, without and after correction for intrinsic stellar extinction (vertical lines connecting symbols at equal  $R^*$ ). The locus in this diagram where the  $Ly_c$  output from the pAGB component precisely accounts for the observed  $H\alpha$  luminosity is depicted by the thick horizontal line at  $\log(\tau)=0$ , and the dashed line at  $\log(\tau)=0.3$  marks the boundary between pAGB photoionization and  $Ly_c$  escape (cf P13).

Panel n: Luminosity contribution ( $\mathcal{L}_{5 \text{ Gyr}} [\%]$ ) of stellar populations younger than  $5$  Gyr at  $5150$  Å (normalization  $\lambda$ ).

In the Appendix D we additionally include five intermediate-to-late type CALIFA galaxies that were processed with PORTO3D in the same manner as ETGs, except for using only BC03 SSPs for spectral fitting. All but one of these systems are classifiable by the BPT ratios in their central part and/or extranuclear component as *composite* or *SF-dominated* (C and H, respectively, in the notation of Table 1; col. 13). These non-ETG galaxies (not further discussed here) are only meant to illustrate how several quantities of interest, and their uncertainties, vary from almost *wim*-devoid type ii ETGs towards SF-dominated late-type galaxies with an  $EW(H\alpha)$  several times larger than  $EW_{\star}^{+}$ . For example, it is interesting to note how the fractional area in which extra (non-pAGB) photoionization needs be postulated (yellow regions in panel f) strikingly increases from ETGs towards “later” galaxy types, reaching in the case of NGC 4470  $\sim 90\%$  of the total spectroscopically studied area. This trend is accompanied by a predominance of stars younger than  $5$  Gyr to the optical luminosity (panel n). Noteworthy is also the reduction of the mean radial  $\log([\text{O III}]/H\beta)$  and  $\log([\text{N II}]/H\alpha)$  ratios – both now offset by  $\gtrsim -0.5$  dex relative to the LINER-typical ratios of ETGs – placing NGC 4470 throughout its extent into the ‘composite/SF’ regime of the BPT plane.



**Fig. 1.** NGC 1167: 2D maps (panels a-f and n), radial intensity and EW of  $\text{H}\alpha$  (panels g and h), BPT diagrams (panels i and j), radial distribution of diagnostic line ratios (k and l), and the  $\tau$  ratio (panel m). In all 2D maps north is up and east to the left. The horizontal bar corresponds to  $20''$ .



## 5. Discussion

### 5.1. 2D properties of gas and stars in the sample ETGs

The  $\mathcal{L}_{5\text{ Gyr}}$  maps (Figs. 1n–C.31n) echo the well established fact that ETGs are dominated by an evolved stellar component throughout their optical extent. Even though there is a trend for an increasing luminosity contribution from stars younger than 5 Gyr towards the galaxy periphery, in agreement with the conclusions by, e.g., [González Delgado et al. \(2014a\)](#), panels n reveal, just like the inwardly increasing mass-weighted stellar age  $\langle t_\star \rangle_M$  (cf Fig. B.3), that the bulk of the stellar component in ETGs has formed several Gyr ago.

Another insight from the  $H\alpha$  and  $\text{EW}(H\alpha)$  maps (panels b&c) is the presence of a diffuse, low-surface brightness *wim* component over virtually the entire optical extent of our sample ETGs. Its co-spatiality with the evolved underlying background is consistent with the notion that the diffuse ionizing field from the latter is an important driver of extranuclear *wim* emission. Following our pilot study in K12, this issue was addressed in more detail and on the basis of the robust statistics from this sample in P13, where we concluded that, in about 40% of ETGs (type i), the properties of the extranuclear *wim* are indeed compatible with the pAGB photoionization hypothesis. The main lines of arguing in those studies were based on a combined analysis of the  $\text{EW}(H\alpha)$  and the  $\tau$  ratio.

Regarding the first quantity, one can read off Fig. A.1 that the photoionization by pAGB stars can reproduce the narrow range of EWs between  $\text{EW}_\star^-$  (0.5 Å) and  $\text{EW}_\star^+$  ( $\sim 2.4$  Å). This range can be further limited to  $\approx 1$  Å ( $\equiv \langle \text{EW}_\star \rangle$ ) when taking into account the typical  $\langle t_\star \rangle_M$  (7 – 11 Gyr) of ETGs, and making the reasonable assumption of a nearly-solar metallicity (see discussion in Sect. A). Obviously, the condition  $\text{EW}_\star^- \leq \text{EW}(H\alpha) \leq \text{EW}_\star^+$  ensures compliance of observations with predictions for pure pAGB models.

In passing we note that we do not invoke the presence or absence of LINER characteristics as an argument for or against pAGB photoionization, since it is known that they can be due to a variety of mechanisms, including shocks and starburst-driven outflows (see, e.g., [Sharp & Bland-Hawthorn, 2010](#), for a recent observational review). For example, the biconical *wim* component protruding several kpc away from the nucleus of NGC 5966 in NE-SW direction (Fig. C.4; see K12 for a detailed discussion) shows throughout LINER-typical emission-line ratios and an  $\text{EW}(H\alpha)$  that is compatible with pure pAGB photoionization. Notwithstanding this fact, the morphology of this kinematically decoupled *wim* component is suggestive of large-scale shock excitation associated to gas outflow. The absence of ongoing SF throughout this ETG lends support to the idea that its gas outflow is powered by an AGN hosted in its LINER nucleus.


As for the  $\tau$  ratio, a radial constancy around unity would ensure that pAGB stars are capable of supplying the ionizing photon budget that is necessary for sustaining the observed  $H\alpha$  emission without need for any additional excitation source. The latter case would correspond to  $\tau < 1$ , whereas a  $\tau > 1$  implies the escape of a fraction  $p_{lf} = 1 - \tau^{-1}$  of the Lyman continuum radiation produced by the pAGB component (Sect. 1). As argued in [Cid Fernandes et al. \(2011\)](#) a relation between the  $\text{EW}(H\alpha)$  and  $\tau^{-1}$  is to be expected, and it was found indeed from an analysis of Sloan Digital Sky Survey (SDSS) data (see their Fig. 5). Likewise, a tight anticorrelation between  $\tau$  and  $\text{EW}(H\alpha)$  over nearly 3 dex, with the pAGB “equilibrium” line of  $\tau \approx 1$  corresponding to  $\text{EW}(H\alpha) \approx 1$  Å ( $\equiv \langle \text{EW}_\star \rangle$ ) was documented from the analysis of CALIFA IFS data in P13 (see their Fig. 2a).

One may superficially argue that  $\text{EW}(H\alpha)$  and  $\tau$  reflect essentially one and the same quantity, they therefore can be used interchangeably in assessing the validity of pAGB photoionization hypothesis. However, an appreciation of the a) geometry-dependent line-of-sight dilution effect of nuclear EWs in triaxial stellar systems and b) the large  $\text{Ly}_c$  escape fraction in the majority of ETGs (see discussion in P13) call this simplistic view into question. For example, a consequence of the geometric dilution is that a radially constant  $\text{EW}(H\alpha)$  in the range between  $\text{EW}_\star^-$  and  $\text{EW}_\star^+$  can only be regarded as a proof for the validity of the pAGB photoionization hypothesis in the specific case of an oblate stellar system seen nearly face-on, or when in a triaxial stellar system the line-emitting *wim* is uniformly mixed with the EW-diluting stellar background. Whenever the *wim* occupies a smaller volume than the stars, the intrinsic  $\text{EW}(H\alpha)$  within a volume element is greater than the observed (projected) value. This cautionary note is essential for a proper evaluation of panels f where regions satisfying the condition  $\text{EW}_\star^- \leq \text{EW}(H\alpha) \leq \text{EW}_\star^+$  are depicted in open-blue color: pAGB photoionization can be regarded as the dominant gas excitation source in these regions only as long as the *wim* is co-spatial with the stars and  $\text{Ly}_c$  escape is throughout negligible. Would any of these (far from self-evident) assumptions be invalid, then an additional excitation source (e.g., AGN or diffuse SF) would need to be postulated. In fact, comparison of panels b & c shows the importance of the line-of-sight EW dilution effect in several of our sample galaxies. NGC 5966 offers again an illustrative example: The biconical  $H\alpha$  emission in this system has its maximum at the peak of the stellar surface density (panel b), whereas the opposite is the case for the  $\text{EW}(H\alpha)$  (panel c). Note that a spatial anti-correlation between emission-line fluxes and EWs has been documented and discussed also in other triaxial systems, such as blue compact dwarf galaxies (see, e.g., [Papaderos et al., 2002](#)).

In P13 we showed that, judging from their  $\text{EW}(H\alpha)$  and  $\tau$  profiles in their extranuclear zones ( $R^\star \geq 4''$ ; blue dots in panels g–m), ETGs span a broad, continuous sequence that can be tentatively grouped in two main classes, based on radial  $\text{EW}(H\alpha)$  profiles (see schematic representation in Fig. 2). Type i ETGs (14 galaxies) are characterized by a nearly constant  $\text{EW}(H\alpha)$  of  $\geq 1$  Å out to their periphery, with the exception of a few systems where the EW shows a steep outer increase to  $\text{EW}(H\alpha) \gg \text{EW}_\star^+$ . Such systems, to which [Gomes et al. \(2015a\)](#) assign the notation i+, are subject of a recent parallel CALIFA study ([Gomes et al., 2015b](#)), which reveals that the EW excess is due to low-level star-forming activity in faint embedded spiral-like features. Type ii ETGs (18 galaxies), on the other hand, display very low ( $\leq 0.5$  Å) EWs in their cores and typically positive EW gradients, approaching an  $\text{EW}(H\alpha) \approx \langle \text{EW}_\star \rangle$  only in their outermost periphery ( $R^\star \gtrsim 2 r_p$ ).

### 5.2. Nebular emission in type i ETGs

With regard to most type i ETGs in our sample, our results are consistent with (yet, recalling our cautionary notes in the previous section, no proof for) the hypothesis that the diffuse ionizing photon field from the pAGB component is the main driver of extranuclear *wim* emission. Panels h in Fig. 1 and Figs. C.1–C.31 show that most isan determinations for type i ETGs satisfy the condition  $\text{EW}_\star^- \leq \text{EW}(H\alpha) \leq \text{EW}_\star^+$  (shaded area in panels f), with most data points populating the EW strip around  $\langle \text{EW}_\star \rangle$ . Consistently with this trend, most type i ETGs show a  $\tau \approx 1$ , with all data points located beneath  $\tau = 2$  (dashed horizontal line in panels m).



**Fig. 2.** Schematic representation of the two main classes of ETGs, as defined in P13. The  $\text{EW}(\text{H}\alpha)$  profiles of type i ETGs (blue color), such as, e.g., NGC 1167, show in their extranuclear component (beyond the radius range depicted by the vertical shaded area; typically  $\sim 2$  kpc in our sample) nearly constant values within the narrow range between  $\text{EW}^-$  and  $\text{EW}^+$  (green-color horizontal strip), with a mean value of typically  $\sim 1$  Å. A few of these systems (labeled i+, blue-dashed curve; e.g., NGC 1349) additionally show a steep  $\text{EW}(\text{H}\alpha)$  increase above  $\text{EW}^+$  in their periphery ( $>1$  effective radius), which, as we discuss in Gomes et al. (2015a); Gomes et al. (2015b) (see also Sect. 5.2), is owing to low-level star-forming activity. By contrast, type ii ETGs (red curve) show a centrally very low ( $\leq 0.5$  Å) mean  $\text{EW}(\text{H}\alpha)$ , increasing then smoothly to  $\geq \text{EW}^-$  at their periphery (e.g., NGC 6411). Regarding their nuclear properties, both ETG types display a considerable diversity, from systems with virtually *wim*-evacuated cores ( $<0.5$  Å) to galaxies hosting a compact central core with an  $\text{EW}(\text{H}\alpha)$  of a few  $\text{EW}^+$ . The profiles in light-blue and light-red color are added to illustrate the spread in the nuclear  $\text{EW}(\text{H}\alpha)$ 's of our sample ETGs.

Regarding the subclass of type i+ ETGs, two of them are morphologically classified as S0 (NGC 1349 and NGC 3106), the third one as SA0 (NGC 1167). As shown in Gomes et al. (2015b), the outer EW-enhanced zone of these systems (panel c) is due to photoionization by young stars over a fraction between  $\sim 7\%$  and  $\sim 47\%$  of the galaxy area. Note that on the radial BPT ratio profiles (i.e. the *isan*-based  $[\text{O III}]/\text{H}\beta$  and  $[\text{N II}]/\text{H}\alpha$  determinations in panels k&l) the outer star-forming zone of type i+ ETGs is clearly separable from the LINER-typical average for ETGs (gray strip) only when producing significant nebular emission ( $\geq 6$  Å; NGC 1349 and NGC 3106; Figs. C.12 and Figs. C.13, respectively), yet marginally traceable in the case of NGC 1167 (Fig. 1) where the  $\text{EW}(\text{H}\alpha)$  ( $\sim 3\text{--}4.5$  Å) is just above  $\text{EW}^+$ . As discussed in Gomes et al. (2015b), the spectroscopic BPT classification of type i+ ETGs is strongly dependent on the spectroscopic aperture used (or, equivalently, on the redshift of such a source). The presence of SF activity in the periphery of the three type i+ ETGs in our sample was noticed in previous studies of the CALIFA collaboration (Sánchez, 2014a), even though no explicit discussion was given to this subject due to their specific scope. For example, the analysis underlying the aforementioned studies has established that the outer  $\text{H}\alpha$  rim in NGC 1349 is composed of several genuine HII regions, that is clumpy  $\text{H}\alpha$  entities containing a young ionizing stellar component with a luminosity fraction  $\geq 20\%$  and showing an  $\text{EW}(\text{H}\alpha) \geq 6$  Å (see, e.g., Sánchez et al., 2014b).

Besides the three ETGs classified type i+, our sample contains a few further systems witnessing the potentially relevant role of SF over an extended radial zone or even galactic scale. In fact, the interpretation of pAGB photoionization dominating

at all radii is challenged by singular or contiguous EW enhancements above  $\langle \text{EW}_* \rangle$  in the extranuclear part of several ETGs. One such example is the S0 galaxy NGC 6081 (Fig. C.6c). This type i ETG shows LINER characteristics in its nucleus and intermediate zones ( $R^* \leq 10''$ ) notwithstanding a contiguous weak ( $\text{EW}(\text{H}\alpha) \leq 2.5$  Å) SF rim all over its southwestern half. Still, *isan* determinations for the  $\text{EW}(\text{H}\alpha)$  and  $\tau$  are consistent with a dominant contribution from pAGB photoionization, in agreement with Fig. C.6f which indicates that sources other than pAGB stars dominate in less 7% of the area of the galaxy. Weak embedded SF patterns are also visible in three intermediate-morphology galaxies, NGC 7025 (Fig. C.11b&c), UGC 10205 (Fig. C.5b&c) and NGC 4003 (Fig. C.2b&c). In particular,  $\text{H}\alpha$  and  $\text{EW}(\text{H}\alpha)$  maps for NGC 7025 reveal a complex, low-level SF network with an  $\text{EW}(\text{H}\alpha) > \text{EW}^+$ , surrounding an extended ( $R^* \sim 10''$ ) LINER/AGN core. LINER characteristics are also apparent in the intermediate zone of NGC 4003, whereas the nuclear region and the periphery of this ETG are dominated by SF. This case also illustrates how the area considered in the spectroscopic analysis can influence the spectroscopic classification, hence the relative role ascribed to different gas excitation mechanisms: *sisp* (i.e. higher-S/N) data capture only the intermediate LINER zone surrounding the SF nucleus, whereas the SF-dominated low-surface brightness periphery of the galaxy becomes apparent on  $\log([\text{O III}]/\text{H}\beta)$  and  $\log([\text{N II}]/\text{H}\alpha)$  profiles only through *isan* determinations including lower-S/N spaxels.

As for the nuclear properties of type i ETGs, all except for NGC 4003 and UGC 10205 fall, according to our classification in Table 1 into the LINER locus of BPT diagrams, just like our ETG sample as a whole. Note that *sisp* determinations on the  $\log([\text{N II}]/\text{H}\alpha)$  vs  $\log([\text{O III}]/\text{H}\beta)$  and  $\log(\Sigma[\text{S II}]/\text{H}\alpha)$  vs  $\log([\text{O III}]/\text{H}\beta)$  diagrams (panels i&j) place the nuclei and intermediate zones close to the junction between the curves which demarcate the regions of ‘maximum SF’ and ‘AGN excitation’.

Our study also shows that type i ETGs span a wide range in the  $\text{EW}(\text{H}\alpha)$  and spatial extent of their nuclear emission. A few of them show prominent nuclear *wim* components over  $\sim 6''\text{--}10''$  (e.g., NGC 1167, NGC 5966, UGC 5771, NGC 6146) with a peak EW of up to  $\sim 8$  Å, whereas in others the nuclear nebular emission is rather compact ( $\sim \text{FWHM}$ ) and faint ( $\sim \langle \text{EW}_* \rangle$ ). One example of extended nuclear emission over scales of  $\sim 4$  kpc is seen in NGC 1167. Besides the biconical *wim* protruding out to  $\sim 10$  kpc from the nucleus of NGC 5966 in a direction perpendicular to the major axis, bipolar or unipolar *wim* lobes on kpc-scales from the nucleus are detected in UGC 10695 and NGC 6146. A robust spectroscopic and kinematical analysis of these extended, faint ( $\text{EW}(\text{H}\alpha) \leq \langle \text{EW}_* \rangle$ ) features is barely possible with the available CALIFA V500 data due to their low spectral and spatial resolution ( $\text{FWHM} \geq 1$  kpc at their distance of the four ETGs), and medium S/N. However, the absence of appreciable star formation in these ETG nuclei renders the interpretation of the detected *wim*lobes as AGN-driven outflows plausible, making them targets of considerable interest for follow-up studies exploring direct evidence for accretion-powered nuclear activity in LINER nuclei. Spatially resolved IFS studies of high-excitation (i.e. AGN-specific) emission lines in the nuclear and circumnuclear region of these candidate AGN (following the approach by Müller-Sánchez et al., 2011), and supplemented by kinematical modeling of biconical gas outflows possibly emanating from their narrow-line regions (e.g., Fischer et al., 2013) could provide key insights in this respect.

### 5.3. The deficit of nebular emission in type ii ETGs

The nature of type ii ETGs – systems commonly referred to as ‘ETGs without nebular emission’ – is enigmatic. These galaxies make about 60% (18 of 32) of our sample and are classified on the condition  $\langle \tau \rangle \geq 2$  in their extranuclear component (cf P13). Nebular emission in these systems is only detectable after accurate fitting and subtraction of the local underlying stellar SED, which clearly is a demanding task, given its extreme faintness. This is particularly true for the virtually *wim*-evacuated cores of these ETGs, where the  $\text{EW}(\text{H}\alpha)$  has its minimum, rising then gradually to levels  $\geq \text{EW}_\star^-$  up to  $\langle \text{EW}_\star \rangle$  in the outermost galaxy periphery ( $\sim 2r_p$ ). These positive  $\text{EW}(\text{H}\alpha)$  gradients, as distinctive feature of type ii ETGs, were interpreted by P13 as manifestation of the radius-dependent structure and physical conditions of the gas in these systems (volume filling factor, porosity and 3D distribution; electron density and ionization parameter). Panels f show that the low  $\text{EW}(\text{H}\alpha)$  within typically more than 80% of the spectroscopically studied area of type ii ETGs is incompatible with pAGB photoionization, if case B recombination for standard gas conditions ( $n_e = 100 \text{ cm}^{-3}$ ,  $T_e = 10^4 \text{ K}$ ) is assumed, a fact underscoring our previous conclusion in P13 that most of the ionizing photons generated by the pAGB component escape without being reprocessed into nebular emission. As a comparison, on average 65% of the area of a type i ETG is consistent with the hypothesis that their gas is being photoionized by pAGB stars.

Diagnostic emission-line ratios and their radial distribution in type ii ETGs have to be considered in the light of their large uncertainties, even after averaging of *sisp* determinations within morphology-adapted irregular annuli (*isan*). Albeit a scatter of typically 0.3 dex for *isan* data, one can consistently infer from panels i&j that type ii ETGs are in practice indistinguishable from type i ETGs by their nearly radially constant LINER-typical BPT ratios.

As for their nuclear properties, type ii ETGs also feature a considerable heterogeneity: From panels b&c it can be seen that in some of these systems a compact yet extended ( $\geq 3$ -PSF) nuclear  $\text{H}\alpha$  excess is robustly detected, as, for example, in NGC 2918, NGC 6150, NGC 6173, NGC 6515, NGC 7550 and, more impressively, in UGC 11958 and NGC 6338. In the majority of type ii ETGs, however, nebular traces of nuclear activity are almost absent, with an  $\text{EW}(\text{H}\alpha)$  marginally above that of the diffuse *wim* and at the edge of the detectability (0.1–0.3 Å).

With regard to their peripheral zones, whereas individual *sisp* data points are generally subject to prohibitively large relative uncertainties (above 100% at  $\text{EW}_\star < 0.3 \text{ Å}$ ), the averaging of hundreds of such determinations within *isan* shows the presence of a faint substrate of *wim* with an  $\text{EW}(\text{H}\alpha) \geq \text{EW}_\star^-$ . This is augmented by determinations of emission-line fluxes based on azimuthally binned *isan* segments (Fig. B.5) or the collapsed spectrum within each *isan*, after correction of local stellar motions (Fig. B.6). It should be noted, on the other hand, that the angular resolution of our data and the faintness of the *wim* does not allow us to determine whether part of the nebular emission originates from higher-density HII-region clumps within a more tenuous *wim* component.

#### 5.3.1. Accretion-powered nuclear activity in ETG/LINERs

Whether or not LINER nuclei in ETGs harbor an AGN is subject of an intense longstanding controversy, with antipodal conclusions reached, including the radical rejection of the hypothesis of accretion-powered nuclear activity in favor of pure pAGB

photoionization (cf Introduction). However, a first reservation towards the latter statement comes from the established relation between the velocity dispersion of galaxy spheroids and the mass of their super-massive black hole (SMBH) (see Merritt & Ferrarese, 2001, for a review), which is estimated to range between a few  $10^7 M_\odot$  and up to  $\sim 10^{10} M_\odot$ . If so, a minimum amount of gas would in principle suffice for sustaining accretion-powered nuclear activity, perhaps in form of a ‘weak’, radiatively inefficient LLAGN. As this study shows, diffuse *wim* is present, even in type ii ETGs, leaving space for the AGN hypothesis. Indirect observational support to the latter comes from the bi-conical/unipolar *wim* lobes in three type i ETGs of our sample (Sect. 5.2), which are consistent with AGN-driven gas outflows. Additionally, in one type i ETG (NGC 1167) the presence of an AGN is indicated by a radio jet (e.g., Sanghera et al., 1995; Giovannini et al., 2001). Also, among type ii ETGs, radio-continuum data for NGC 6150 and UGC 11958 (cf Table 1) witness the energy release by an AGN. For example, the line-weak ( $\text{EW}(\text{H}\alpha) \sim 3.5 \text{ Å}$ ) LINER nucleus of UGC 11958 powers a  $\geq 100$  kpc double-lobe radio continuum halo detected at 1.4 GHz with the VLA (Comins & Owen, 1991) with a large ( $\sim 40$  kpc) central cavity of hot ( $\sim 1 \text{ keV}$ ), pressure-driven X-ray emitting plasma (Worrall et al., 2007), clearly indicating the energetic action by an AGN.

How to reconcile the weakness of optical emission lines in ETG/LINERs with signatures of strong accretion-powered nuclear activity in radio- and X-ray wavelengths poses an enigma. As first pointed out in P13, the seemingly trivial fact that more than one half of ETGs lacks nebular emission as a result of an insufficient mean gas density and/or high gas porosity has far-reaching consequences for our understanding of accretion-powered nuclear activity in these systems: In the presence of extensive  $\text{Ly}_\gamma$  photon escape, implied by the lack of gas, emission-line luminosities and EWs, upon which estimates of accretion rates and SMBH masses rely, are reduced by more than an order of magnitude. An additional effect acting towards diminishing nuclear EWs (or any photometric signs of nuclear activity) in these triaxial stellar systems is dilution by the stellar background along the line of sight (P13). According to our study in P13, the line weakness of ETG/LINER nuclei is therefore no foolproof evidence for the weakness or absence of AGN in these systems. In the light of this conclusion and from the evidence presented in Sects. 5.2&5.3, it is therefore conceivable that different gas excitation mechanisms (photoionization by AGN, OB stars and pAGB sources; shocks) – each of them with a different relative contribution in different radial zones – are simultaneously at work. An exploration of this issue with deep, higher-spectral resolution IFS data is apparently of great interest.

### 5.4. Stellar and gas kinematics in ETGs

A detailed study of stellar and gas kinematics in the CALIFA galaxy survey is subject of several ongoing investigations within the CALIFA collaboration (García-Lorenzo et al., 2015; Sánchez-Blázquez et al., 2014, Falcón-Barroso et al., in prep.), which use higher-resolution (V1200; FWHM  $\sim 2.3 \text{ Å}$ ) IFS data and to which the reader is referred for detailed information. The brief remarks below are merely meant to further highlight the kinematical diversity of our sample ETGs.

As already revealed by several studies before, most notably in the SAURON and Atlas<sup>3D</sup> projects (cf, e.g., Sarzi et al., 2006, 2010), ETGs display a variety in their stellar kinematics patterns, from regular rotation along the semi-major axis to



nearly pressure-supported galaxies with subtle, if any, signs of ordered motions.

Panels d&e of Fig. 1–C.31 show that in most (27) of the ETGs studied here stellar kinematics are dominated by rotation. The velocity difference  $\Delta v_{\star} = (v_a + v_r)/2$  between the receding ( $v_r$ ) and approaching ( $v_a$ ) part of these systems is typically of the order of  $200 \text{ km s}^{-1}$ . Interestingly, weak rotation at  $60 \text{ km s}^{-1} \lesssim \Delta v_{\star} \lesssim 110 \text{ km s}^{-1}$  along the minor galaxy axis is clearly detected in three type ii ETGs (NGC 6173, NGC 6338 and NGC 7236). Five galaxies (UGC 10695, NGC 6515, UGC 11958, NGC 7550 and UGC 29) appear to be dominated by random motions. Note, however, that the low spectral resolution and medium S/N of our CALIFA V500 IFS data, do not permit to firmly rule out some degree of rotation in these systems. Higher-spectral resolution CALIFA data (Falcón-Barroso et al., in prep.) enabling a detailed study of stellar line-of-sight velocity patterns and possible kinematical distortions (e.g., counter-rotating cores) over the full set of CALIFA ETGs are necessary for a conclusive assessment of this issue. An interesting question in this respect is how type i/i+ and type ii ETGs compare to each other with respect to their angular momentum per unit mass in the definition by Emsellem et al. (2007). These authors have shown that fast rotators in the SAURON sample tend to be lower-luminosity ETGs with well-aligned photometric and kinematic axes, whereas slow rotators typically exhibit significant misalignments and kinematically decoupled cores.

A comparative study of type i/i+ and ii ETGs with regard to the kinematics of their *wim* is obviously another key task that could help understanding whether gas is internally produced, in the course of stellar evolution, or accreted from captured satellites. The spiral-arm like SF features in type i+ ETGs, or faint embedded SF patterns in several other ETGs are suggestive, yet no proof, for the gas accretion scenario. With regard to the discussion below it is important to bear in mind that the non-detection of ordered gas motions in most of our sample ETGs (especially those classified type ii) may reflect observational limitations due to the extreme faintness of the *wim* (see also Sect. 5.5, in particular our remarks on the type i+ ETG NGC 160 in Fig. 3). Clearly, a detailed study of ionized-gas kinematics is, in particular in type ii ETGs a formidable task that only can be pursued with a high S/N (typically  $\geq 60$  at  $6390\text{--}6490 \text{ \AA}$ ) and/or spatially binned data (e.g., García-Lorenzo et al., 2015).

Some noteworthy trends are apparent though from the analysis of our low-resolution V500 data: In roughly one third (11) of the sample ETGs (8 classified type i and in the three type i+) gas kinematics are consistent with rotation aligned with stellar motions, albeit slightly perturbed kinematical patterns. For instance, in NGC 1167 the *wim* rotation axis appears slightly tilted relative to the stellar one, with a possible spatial offset between the photometric and kinematical center. In three type i ETGs (NGC 5966, UGC 10695 and UGC 10905) gas kinematics appear to be dominated by outflows, and a suggestion for an outflowing gas component is also present in NGC 6146. Of the three type ii ETGs showing stellar rotation perpendicular to the major axis, we do not detect gas motions in NGC 6173 and NGC 7236, whereas in NGC 6338 there is a trace of a compact rotating ( $\sim 70 \text{ km s}^{-1}$ ) *wim* component, centered on the optical nucleus. Likewise, in NGC 7550 gas rotation at  $\Delta v_g \sim 80 \text{ km s}^{-1}$  can be traced within the central  $\sim 7 \text{ kpc}$  of a pressure-supported stellar host. Summarizing, this study further highlights and adds observational insight into the kinematical diversity of the ETG galaxy family (see also, e.g., Sarzi et al., 2006; Katkov et al., 2014). On the other hand, the evidence from Figs. 1 and C.1–C.31 permits identification of broad kinematical trends in the three ETG


types: Type i and i+ ETGs are generally rotation-supported with many cases of perturbed rotational patterns or outflows in their *wim*, whereas type ii ETGs exhibit both rotationally or pressure supported stellar kinematics, and generally lack distinguishable kinematic patterns in their ionized gas. In Table 1 (col. 14) includes a descriptive 3-character kinematical classification, with **r** denoting rotation-dominated kinematics, **p** perturbed rotational patterns, **o** gas outflows, and **n** no distinguishable patterns or pressure-supported stellar kinematics. The first and third character refer, respectively, to the kinematics of stars and gas, and the second one indicates in the case of stellar rotation (r or p) whether the kinematic and photometric major axes are roughly aligned (||) or not (⊥).

### 5.5. Equivalent width bias on ETG taxonomy

A central result from our study in P13 is the considerable radial dependence of the  $\text{EW}(\text{H}\alpha)$  in many ETGs (panels c&h): A distinctive property of several type ii ETGs is their almost *wim*-emission free cores and their outwardly increasing  $\text{EW}(\text{H}\alpha)$ , reaching in some cases readily detectable levels ( $\sim \langle \text{EW}_{\star} \rangle$ ) in their periphery only. With regard to type i ETGs, although these show on average an  $\text{EW}(\text{H}\alpha) \approx \langle \text{EW}_{\star} \rangle$  in their extranuclear component, there are several cases in our sample where the  $\text{EW}(\text{H}\alpha)$  significantly varies within different radial zones. This is particularly true for the two-zone *wim* morphology of type i+ ETGs: An illustrative example NGC 1349 showing a constant  $\text{EW}(\text{H}\alpha) \approx \langle \text{EW}_{\star} \rangle$  for  $R^* \leq 8''$  (3.4 kpc), in agreement with pure pAGB photoionization, whereas in the periphery ( $R^* \sim 16\text{--}20''$ ) the  $\text{EW}(\text{H}\alpha)$  rises to  $\geq 6 \text{ \AA}$ , revealing an extra ionizing field produced by ongoing SF activity (Sect. 5.2).

In fact, the intuitive expectation that emission-line EWs in ETGs are spatially correlated with the surface brightness of the stellar background, invariably attaining their peak value at the ETG nuclei, is disproved by panels c and h of Figs. 1–C.31. What a comparison of panels a&b with panel c reveals in many cases is actually an inverse trend with the  $\text{EW}(\text{H}\alpha)$  showing – contrary to the stellar and  $\text{H}\alpha$  surface brightness – low or even minimum value in the nuclear region (e.g., Fig. C.4). This spatial anti-correlation between the intensity and EW of emission lines has been documented and discussed in studies of low-mass starburst galaxies (Papaderos et al., 1998, 2002) and, in the case of ETGs, it can partly attributed to line-of-sight dilution of nuclear EWs by a triaxial stellar host (P13). As we pointed out in P13, this geometric effect introduces a selection bias against the detection of nuclear activity in galaxy spheroids, since emission-line EWs (and obviously any other manifestation of energy release by an AGN in UV-NIR wavelengths) in their centers are diluted, not only by the local stellar background, but, additionally, by the integrated line-of-sight luminosity of stars. Since this geometric dilution effect amplifies an observational bias stemming from the high  $L_{\text{yc}}$  photon escape fraction in type ii ETG nuclei, caution is needed when arguing that the weakness of nuclear emission-lines and EWs in ETGs/LINERs is evidence for weak or non-existent accretion-powered nuclear activity.

In the light of these two effects, analogous considerations apply also to the usage of  $\text{EW}(\text{H}\alpha)$  as discriminator between AGN and ‘retired’ galaxies in the context of the WHAN classification (Cid Fernandes et al., 2011): A nuclear  $\text{EW}(\text{H}\alpha)$  between  $\text{EW}_{\star}^{-}$  and  $\text{EW}_{\star}^{+}$ , i.e. in the range of predicted values for pAGB photoionization in ‘retired’ galaxies, does not necessarily rule out AGN as the dominant excitation source of the *wim*. The true nuclear EW (i.e. the EW after correction for the geo-



**Fig. 3.** Emission-line free stellar continuum (a),  $H\alpha$  flux in  $10^{-16} \text{ erg s}^{-1} \text{ cm}^{-2}$  (b) and  $H\alpha + [\text{NII}]$  line-of-sight velocity in km/s (c) of the type i+ ETG NGC 160, as inferred from processing of CALIFA (v1.5) V500 with PORTO3D. The overlaid contours delineate the  $\text{EW}(H\alpha)$  morphology of the galaxy and go from  $4 \text{ \AA}$  to  $21 \text{ \AA}$  in steps of  $3 \text{ \AA}$ . North is up and east to the left.

metric dilution and  $L_{\text{y}}$  escape bias) could exceed by more than one order of magnitude the observed (projected) one, shifting on the WHAN diagram a seemingly ‘weak’ (low-luminosity) AGN into the ‘strong’ AGN regime (cf our discussion in P13).

It is also worth contemplating the observational biases that the EW detection limit may have on taxonomical studies of ETGs. Note that extranuclear nebular emission would not had been detected in any type ii ETG if our analysis was restricted to an  $\text{EW}(H\alpha) \geq 1 \text{ \AA}$ . Because of the geometrical EW dilution effect, even the compact  $H\alpha$ -emitting nuclei of many type i and type ii ETGs (e.g., in NGC 1349, UGC 8234, NGC 6515, NGC 7025) vanish to  $\text{EWs} \leq 1 \text{ \AA}$ , and would had gone undetected if such a ‘quality-ensuring’  $\text{EW}(H\alpha)$  threshold would had been adopted. By that cutoff, all but three type ii ETGs (NGC 6338, UGC 11958 and NGC 7550) would had been classified as gas-free ‘passive’ galaxies (see definition by Cid Fernandes et al., 2011). Even though the thin higher-EW outer rim of nebular emission would had been recovered in type i+ ETGs, this alone would not had demonstrated the extended nature of the circumnuclear *wim* component.

The importance of this EW-threshold bias may be illustrated through inspection of the *wim* morphology and kinematics maps of NGC 160, a type i+ ETG which will be subject of a forthcoming article of this series. The overlaid contours in the three panels of Fig. 3 depict the  $\text{EW}(H\alpha)$  morphology of the galaxy and go from  $4 \text{ \AA}$  to  $21 \text{ \AA}$  in steps of  $3 \text{ \AA}$ . Whereas a substrate of diffuse ( $0.5 \leq \text{EW}(H\alpha) [\text{\AA}] \leq 2$ ) *wim* is present almost everywhere in the central part of the galaxy, the bulk of  $H\alpha$  emission is confined to an extended ( $\sim 60'' \times 40''$ ) rim surrounding the low-surface brightness stellar periphery (see also García-Lorenzo et al., 2015). An EW detection threshold of  $4 \text{ \AA}$  (i.e.  $\geq \text{EW}_{\star}^{+}$ ; outermost contour in all panels) would obviously allow for the recovery of the outer rim only, leading to the erroneous conclusion that this ETG is completely devoid of *wim* almost throughout its optical extent, out to  $\sim 24 \text{ mag}/\square''$  in the SDSS *r* band (cf Breda et al., in prep.). A further implication would had been the complete rejection of the pAGB photoionization hypothesis, since this is testable only for an EW detection threshold  $\leq \text{EW}_{\star}^{+}$ , and the conclusion that photoionization by OB stars is the sole gas excitation mechanism in this ETG. An EW threshold  $\geq \text{EW}_{\star}^{+}$  would also result in a severely incomplete view of the overall gas kinematics everywhere but in the EW-enhanced rim, with obvious implications to the dynamical modeling of such a galaxy.

## 6. Summary and conclusions

In this article, we investigate in significantly more detail the sample of 32 CALIFA ETGs that we discussed in P13 which had its focus on their radial  $\text{EW}(H\alpha)$  distribution and  $L_{\text{y}}$  photon escape fraction. We study here the 2D properties of their diffuse warm interstellar medium (*wim*) to gain insight into its excitation mechanism(s) and provide additional observational constraints aiming at systematizing the physical properties of these systems. Among other quantities derived by applying our IFS data processing pipeline PORTO3D to low-spectral resolution ( $\sim 6 \text{ \AA}$  FWHM) CALIFA data, we present for each galaxy  $H\alpha$  intensity and EW maps, as well as radial profiles. Also, we analyze the nuclear and extranuclear nebular emission of the *wim* using diagnostic line ratios. A comparison sample of intermediate-to-late type CALIFA galaxies is meant to illustrate how various observables of interest change from bulge-dominated ETGs towards disk-dominated star-forming (SF) galaxies. Additionally, we briefly discuss the kinematical properties of gas and stars in our sample.

The main conclusions from this study may be summarized as follows:

i) The analyzed ETGs contain an extended *wim* component displaying a considerable diversity in its mean  $\text{EW}(H\alpha)$  and its radial gradients. As we discussed in P13, our ETG sample can be tentatively subdivided in two groups. The extranuclear *wim* of the first class (type i) is characterized by a nearly constant  $\text{EW}(H\alpha)$  of  $\sim 1 \text{ \AA}$ , with the exception of a few systems (i+, in the notation by Gomes et al., 2015b) showing a steep  $\text{EW}(H\alpha)$  increase in their periphery. The second group (type ii) displays almost *wim*-evacuated cores and a gradual increase of the  $\text{EW}(H\alpha)$  outwardly to  $\lesssim 1 \text{ \AA}$ .

ii) The ETGs studied here also display a large diversity in their kinematics. The majority (27 out of 32) of our sample ETGs exhibit stellar kinematics dominated by rotation, with three systems showing a kinematic major axis aligned with the photometric minor axis. Type i and i+ ETGs are generally rotation-supported with many cases of perturbed rotational patterns or outflows in their ionized gas, whereas the class of type ii ETGs contains both rotationally and pressure supported stellar systems, which generally lack distinguishable kinematic patterns in their ionized gas.

iii) We compute the time evolution of the  $\text{EW}(H\alpha)$  for various star formation histories (SFHs) and stellar metallicities based on an evolutionary synthesis model and assuming case B recombination and standard conditions for the gas. We find that SFHs

involving an exponentially decreasing star formation rate since 13 Gyr can reproduce the low ( $\sim 1 \text{ \AA}$ )  $\text{EW}(\text{H}\alpha)$ 's in the extranuclear component of present-day ETGs when a short ( $\leq 1 \text{ Gyr}$ ) e-folding timescale is adopted. In this case, the Lyman continuum ( $L_{\text{yc}}$ ) output predicted to arise from the pAGB stellar component is sufficient for powering the excitation of the diffuse *wim*. SFHs involving a longer e-folding timescale (e.g.,  $\geq 3 \text{ Gyr}$ ) imply higher  $\text{EW}(\text{H}\alpha)$ 's than observed ones.

iv) As far as most type i ETGs are concerned, photoionization by the evolved pAGB stellar background offers a consistent, yet not necessarily compelling, explanation for the observed  $\text{EW}(\text{H}\alpha)$ 's and  $\tau$  ratios (the inverse ratio of the observed  $\text{H}\alpha$  luminosity to that predicted for pure pAGB photoionization). We point out that  $L_{\text{yc}}$  photon escape and geometric line-of-sight dilution of EWs (cf P13) could conspire in such a way as to reproduce the low ( $\approx 1 \text{ \AA}$ ) and nearly constant  $\text{EW}(\text{H}\alpha)$  in the extranuclear component of type i ETGs, thereby mimicking a predominance of pAGB photoionization to the excitation of their diffuse ionized gas component, while hiding a potentially important role by AGN and low-level SF. The presence of the latter in the extranuclear component of several ETGs from our sample is clearly indicated from  $\text{EW}(\text{H}\alpha)$  maps.

v) The radial  $\text{EW}(\text{H}\alpha)$  gradients and the extreme faintness of nebular emission in ETGs, in particular in systems classified type ii, leads to biases impacting the detectability of the extended *wim* component and the interpretation of its excitation mechanisms. We point out that nebular emission in the centers of type ii ETGs will generally evade detection below an EW threshold of  $\sim 0.5\text{--}1.0 \text{ \AA}$ , which could lead to the conclusion that these systems are entirely devoid of *wim*, and their erroneous classification as passive galaxies.

vi) The radial distribution of diagnostic line ratios indicates that, in their majority, ETGs show LINER-specific spectroscopic properties both in their nuclear and extranuclear zones. Our sample includes, however, a few notable exceptions (in particular, type i+ ETGs) where a significant contribution by star formation in peripheral zones is reflected both on radial  $\log([\text{N II}]/\text{H}\alpha)$ ,  $\log([\text{O III}]/\text{H}\beta)$  and  $\tau$  profiles, and  $\text{EW}(\text{H}\alpha)$  maps. This calls attention to the heterogeneity of the ETG class, pointing to a complex interplay between the 3D properties of the *wim* and the combined action of various gas ionization sources, such as, e.g., shocks, OB stars and the evolved pAGB component. As we argue in Papaderos et al. (2013) and further detail here, the weakness of nebular emission lines in ETG nuclei is not compelling evidence for the absence of accretion-powered nuclear activity in these systems. In fact, radio-continuum and/or X-ray data, or bipolar or unipolar *wim*lobes protruding several kpc away from the LINER nuclei of 25% of our sample ETGs witness the energetic action of an AGN. Summarizing, the results and consideration presented in this study are incompatible with the hypothesis of the global gas excitation patterns in ETGs being controlled by one single dominant mechanism.

**Acknowledgements.** This paper is based on data from the Calar Alto Legacy Integral Field Area Survey, CALIFA (<http://califa.caha.es>), funded by the Spanish Ministry of Science under grant ICTS-2009-10, and the Centro Astronómico Hispano-Alemán. JMG acknowledges support by Fundação para a Ciência e a Tecnologia (FCT) through the Fellowship SFRH/BPD/66958/2009 and POPH/FSE (EC) by FEDER funding through the program Programa Operacional de Factores de Competitividade (COMPETE). PP is supported by FCT through the Investigador FCT Contract No. IF/01220/2013 and POPH/FSE (EC) by FEDER funding through the program COMPETE. JMG&PP also acknowledge support by FCT under project FCOMP-01-0124-FEDER-029170 (Reference FCT PTDC/FIS-AST/3214/2012), funded by FCT-MEC (PIDDAC) and FEDER (COMPETE). SFS acknowledges support from CONACyT-180125 and PAPIIT-IA100815 grants. Support for LG is provided by the Ministry of

Economy, Development, and Tourism's Millennium Science Initiative through grant IC120009, awarded to The Millennium Institute of Astrophysics, MAS. LG acknowledges support by CONICYT through FONDECYT grant 3140566. CJW acknowledges support through the Marie Curie Career Integration Grant 303912. RGB is supported by the Spanish Ministerio de Ciencia e Innovación under grant AYA2010-15081. R.A. Marino is funded by the Spanish program of International Campus of Excellence Moncloa (CEI). IM acknowledges financial support by the Junta de Andalucía through project TIC114, and the Spanish Ministry of Economy and Competitiveness (MINECO) through projects AYA2010-15169 and AYA2013-42227-P. AdO acknowledges financial support from the Spanish grants AYA2010-15169 and AYA2013-42227-P. J. F-B acknowledges support from grant AYA2013-48226-C3-1-P from the Spanish Ministry of Economy and Competitiveness (MINECO). The STARLIGHT project is supported by the Brazilian agencies CNPq, CAPES, and FAPESP. We benefited from stimulating discussions with several members of the CALIFA collaboration. This research made use of the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration.



## References

- Allen, M. G., Groves, B. A., Dopita, M. A., Sutherland, R. S., & Kewley, L. J. 2008, *ApJS*, 178, 20
- Alongi M., Bertelli G., Bressan A., et al., 1993, *A&AS*, 97, 851
- Annibali, F., Bressan, A., Rampazzo, R., et al. 2010, *A&A*, 519, A40
- Arnold, J. A., Romanowsky, A. J., Brodie, J. P. et al., 2014, *ApJ*, 791, 80A
- Bacon, R., Copin, Y., Monnet, G. et al., 2001, *MNRAS*, 326, 23
- Baldwin, J. A., Phillips, M. M., & Terlevich, R. 1981, *PASP*, 93, 5
- Barrera-Ballesteros, J. K., Falcón-Barroso, J., García-Lorenzo, B., et al. 2014, *A&A*, 568A, 70B
- Bender, R., Surma, P., Döbereiner, S., Möllenhoff, C., & Madejsky, R. 1989, *A&A*, 217, 35
- Binette, L., Magris, C. G., Stasińska, G., & Bruzual, A. G. 1994, *A&A*, 292, 13
- Bregman, J. N. 1978, *ApJ*, 224, 768
- Bressan A., Fagotto F., Bertelli G., Chiosi C., 1993, *A&AS*, 100, 647
- Bruzual, G. & Charlot, S. 2003, *MNRAS*, 344, 1000
- Cappellari, M., Emsellem, E., Krajnović, D. et al. 2011, *MNRAS*, 413, 813
- Castellanos, M., Díaz, A. I., & Terlevich, E. 2002, *MNRAS*, 329, 315
- Chabrier G., 2003, *PASP*, 115, 763
- Cid Fernandes, R., Mateus, A., Sodré, L., Stasińska, G., Gomes, J. M. 2005, *MNRAS*, 358, 363
- Cid Fernandes, R., Asari, N. V., Sodré, L., et al. 2007, *MNRAS*, 375, L16
- Cid Fernandes, R., Stasinka, G., Schlickmann, M. S. et al. 2010, *MNRAS*, 403, 1036
- Cid Fernandes, R., Stasińska, G., Mateus, A. et al. 2011, *MNRAS*, 413, 1687
- Cid Fernandes, R., González Delgado, R. M., García-Benito, R. et al. 2014, *A&A*, 561A, 130C
- Clemens, M. S., Bressan, A., Panuzzo, P., et al. 2009b, *MNRAS*, 392, 982
- Comins, N.F. & Owen, F.N. 1991, *ApJ*, 382, 108
- Demoulin-Ulrich, M., Butcher, H. R., & Boksenberg, A. 1984, *ApJ*, 285, 527
- di Seregho Alighieri, S., Trinchieri, G. & Brocato, E. 1990, in *Windows on Galaxies*, Fabbiano G., Gallagher J. S., Renzini A., eds. *Astrophys. Space Sci. Libr.* Vol. 160, Kluwer, Dordrecht, p. 301
- Dopita, M. A. & Sutherland, R. S. 1995, *ApJ*, 455, 468
- Emsellem, E., Cappellari, M., Krajnović, D. et al. 2007, *MNRAS*, 379, 410
- Eracleous, M., Hwang, J. A. & Flohic, H. M. L. G. 2010, *ApJ*, 711, 796
- Fagotto F., Bressan A., Bertelli G., Chiosi C., 1994a, *A&AS*, 104, 365
- Fagotto F., Bressan A., Bertelli G., Chiosi C., 1994b, *A&AS*, 105, 29
- Fagotto F., Bressan A., Bertelli G., Chiosi C., 1994c, *A&AS*, 105, 39
- Falcón-Barroso, J., Sánchez-Blázquez, P., Vazdekis, A. et al. 2011, *A&A*, 532, 95
- Finkelman, I., Brosch, N., Funes, J. G., Kniazev, A. Y., & Väisänen, P. 2010, *MNRAS*, 407, 2475
- Fischer, T.C., Crenshaw, D. M., Kraemer, S. B., Schmitt, H. R. 2013, *ApJS*, 209, 1
- García-Benito, R., Zibetti, S., Sánchez, S. F., et al. 2015, *A&A*, 576A, 135G
- García-Lorenzo, B., Márquez, I., Barrera-Ballesteros, J. K., et al. 2015, *A&A*, 573A, 59G
- Giovannini, G., Cotton, W.D., Feretti, L. et al. 2001, *ApJ* 552, 508
- Girardi, L., Bressan, A., Chiosi, C., Bertelli, G., & Nasi, E. 1996, *A&AS*, 117, 113
- Gomes, J.M., Papaderos, P., Kehrig, C. et al. 2015a, in *Galaxies in 3D across the Universe*, ed. Ziegler et al., 105
- Gomes, J.M., Papaderos, P., Vílchez, J.M. et al. 2015b, *A&A Letters*, submitted
- González-Delgado, R. M., Pérez, E., Díaz, A. et al. 1995, *ApJ*, 439, 604
- González Delgado, R. M., Pérez, E., Cid Fernandes, R. et al. 2014, *A&A*, 562, A47
- González Delgado, Cid Fernandes, García-Benito et al. 2014, *ApJL*, 791, 16
- Heckmann, T. M. 1980, *A&A*, 87, 152
- Ho, L.C., Filippenko, A. V., & Sargent, W. L. 1995, *ApJS*, 98, 477
- Ho, L. C. 1999, *Adv. Space Res.*, 23, 813
- Ho, L. C. 2008, *ARA&A*, 46, 475
- Husemann, B., Jahnke, K., Sánchez, S. F., et al. 2013, *A&A*, 549, A87H
- Iglesias-Páramo, J., Vílchez, J. M., Galbany, L., et al. 2013, *A&A*, 553, L7
- Katkov, I. Y., Silchenko, O. K. & Afanasiev, V. L. 2014, *MNRAS*, 438, 2798
- Kauffmann, G., Heckman, T. M., Tremonti, C., et al. 2003, *MNRAS*, 346, 1055
- Kaviraj, S., Schawinski, K., Devriendt, J. E. G., et al. 2007, *ApJS*, 173, 619
- Kaviraj, S., Khochfar, S., Schawinski, K., et al. 2008, *MNRAS*, 388, 67
- Kehrig, C., Monreal-Ibero, A., Papaderos, P., et al. 2012, *A&A*, 540, A11 (K12)
- Kelz, A., Verheijen, M. A. W., Roth, M. M., et al. 2006, *PASP*, 118, 129
- Kewley, L. J., Dopita, M. A., Sutherland, R. S., Heisler, C. A., Trevena, J. 2001, *ApJ*, 556, 121
- Kim, D. 1989, *ApJ*, 346, 653
- Kormendy, J., Fisher, D. B., Cornell, M. E., & Bender, R. 2009, *ApJ*, 182, 216
- Kotulla, R., Fritze, U., Weilbacher, P., Anders, P. 2009, *MNRAS*, 396, 462
- Krajnović, D., Karick, A. M., Davies, R. L., et al. 2013, *MNRAS*, 433, 2812K
- Kulkarni S. R., Ofek E. O., Neill J. D., Zheng Z., Juric, M. 2014, *ApJ*, 797, 70K
- Le Borgne, J.-F., Bruzual, G., Pelló, R. et al. 2003, *A&A*, 402, 433L
- Leitherer, C., Schaerer, D., Goldader, J. D., et al. 1999, *ApJ*, 123, 3
- Macchetto, F., Pastoriza, M., Caon, N. et al. 1996, *A&AS*, 120, 463
- Marino, R. A., Rosales-Ortega, F. F., Sánchez, S. F., et al. 2013, *A&A*, 559, A114M
- Mast, D., Rosales-Ortega, F. F., Sánchez, S. F., et al. 2014, *A&A*, 561, A129
- Mathews, W. G. & Baker, J. C. 1971, *ApJ*, 170, 241
- Merritt, D. & Ferrarese, L. 2001, *ApJ*, 547, 140
- Müller-Sánchez, F., Prieto, M. A., Hicks, E. K. S., et al. 2011, *ApJ*, 739, 69
- Noeske, K. G., Papaderos, P., Cairos, L. M., & Fricke, K. J. 2003, *A&A*, 410, 48
- Noeske, K. G., Koo, D. C., Phillips, A. C., et al. 2006, *ApJ*, 640, 143
- Papaderos, P., Izotov, Y. I., Fricke, K. J., Thuan, T. X., Guseva, N. G. 1998, *A&A*, 338, 43
- Papaderos, P., Izotov, Y. I., Thuan, T. X., et al. 2002 *A&A*, 393, 461
- Papaderos, P., Gomes, J. M., Vílchez, J. M., et al. 2013, *A&A*, 555, L1
- Pérez, E., Cid Fernandes, R., González Delgado, R. M., et al. 2013, *ApJ*, 764, L1P
- Phillips, M. M., Jenkins, C. R., Dopita, M. A., et al. 1986, *AJ*, 91, 1062
- Rampazzo, R., Marino, A., Tantaló, R., et al. 2007, *MNRAS*, 381, 245
- Roth, M. M., Kelz, A., Fechner, T., et al. 2005, *PASP*, 117, 620
- Sánchez, S. F., Kennicutt, R. C., Gil de Paz, A., et al. 2012, *A&A*, 538, A8
- Sánchez, S. F., Rosales-Ortega, F. F., Jungwiert, B. et al. 2013, *A&A*, 554, A58S
- Sánchez, S., *Advances in Astronomy, Issue: Metals in 3D: A Cosmic View from Integral Field Spectroscopy*, 2014a, Article ID 596501
- Sánchez, S. F., Rosales-Ortega, F. F., Iglesias-Páramo, J., et al. 2014b, *A&A*, 563, A49
- Sánchez-Blázquez, P., Peletier, R. F., Jiménez-Vicente, J., et al. 2006, *MNRAS*, 371, 703
- Sánchez-Blázquez, P., Rosales-Ortega, F., Méndez-Abreu, J., et al. 2014, *A&A*, 570, A6
- Sanghera, H.S., Saikia, D.J., Luedke, E. et al. 1995, *A&A*, 295, 629
- Sarzi, M., Falcón-Barroso, J., Davies, R. L., et al. 2006, *MNRAS*, 366, 1151
- Sarzi, M., Bacon, R., Cappellari, M., et al. 2007, *NewAR*, 51, 18S
- Sarzi, M., Shields, J. C., Schawinski, K., et al. 2010, *MNRAS*, 402, 2187
- Sarzi, M., Alatalo, K., Blitz, L., et al. 2012, *MNRAS*, in press
- Schawinski, K., Thomas, D., Sarzi, M., et al. 2007, *MNRAS*, 382, 1415
- Schweizer, F. & Seitzer, P. 1988, *ApJ*, 328, 88S
- Shapiro, K. L., Falcón-Barroso, J., van de Ven, G., et al. 2010, *MNRAS*, 402, 2140
- Sharp, R. G. & Bland-Hawthorn, J. 2010, *ApJ*, 711, 818
- Singh, R., van de Ven, G., Jahnke, K., et al. 2013, *A&A*, 558A, 43S
- Sodré, Jr., L. & Stasińska, G. 1999, *A&A*, 345, 391
- Stasińska, G., Vale Asari, N., Cid Fernandes, R., et al. 2008, *MNRAS*, 391, L29
- Stasińska, G., Costa-Duarte, M.V., Vale Asari, N., Cid Fernandes, R., Sodré Jr., L. 2015, *MNRAS*, 449, 559S
- Struck 1999, *Physics Reports*, 321, 1
- Thomas, D., Maraston, C., Schawinski, K., Sarzi, M., Silk, J. 2010, *MNRAS*, 404, 1775
- Trager, S. C., Faber, S. M., Worthey, G., González, J. J. 2000, *AJ*, 120, 165
- Trinchieri, G. & di Serego Alighieri, S. 1991, *AJ*, 101, 1647
- Vazdekis, A., Sánchez-Blázquez, P., Falcón-Barroso, J., et al. 2010, *MNRAS*, 404, 1639
- Veilleux, S. & Osterbrock, D.E. 1987, *ApJS*, 63, 295
- Walcher, J., Böker, T., Charlot, S., et al. 2006, *ApJ*, 649, 692
- Walcher, J., Wisotzki, L., Bekeraité, S., et al. 2014, in prep.
- Weilbacher, P. & Fritze von Alvensleben, U. 2001, *A&A*, 373, L9
- Westera, P., Samland, M., Bruzual, G. & Buser, R. 2002 *A&A*, 381, 524
- White, III, R. E. & Chevalier, R. A. 1983, *ApJ*, 275, 69
- Worrall, D.M., Birkinshaw, M., Kraft, R.P., Hardcastle, M.J. 2007, *ApJ*, 658, L79
- Yan, R. & Blanton, M. R. 2012, *ApJ*, 747:61
- York, D. G., Adelman, J., Anderson, Jr., J. E., et al. 2000, *AJ*, 120, 1579


## Appendix A: The time evolution of the $H\alpha$ equivalent width in the case of pure pAGB photoionization

As discussed in Sect. 1, the time evolution of the  $Ly_c$  output from the evolved pAGB stellar component in ETGs has been evaluated in several previous studies (e.g., Binette et al., 1994; Stasińska et al., 2008; Cid Fernandes et al., 2010) all of which assume an instantaneous SF process. Here we extend this theoretical work by addressing how the  $Ly_c$  rate from the pAGB stellar component, and the associated nebular emission by photoionized gas, evolve when a slowly declining SF activity is assumed. For this purpose, we use our evolutionary synthesis code, adopting a wider range of star formation histories (SFHs), from an instantaneous burst to continuous SF to compute synthetic spectral energy distributions (SEDs).

The synthetic composite stellar SEDs were computed using the full set of ages for the SSPs from Bruzual & Charlot (2003, hereafter BC03; comprising 221 spectra spanning an age between 0 and 20 Gyr) and assuming a constant metallicity  $Z_\odot$ . The BC03 SSP library uses ‘Padova 1994’ evolutionary tracks (Alongi et al., 1993; Bressan et al., 1993; Fagotto et al., 1994a,b,c; Girardi et al., 1996), the Chabrier (2003) IMF with a lower and upper mass cutoff of 0.1 and 100  $M_\odot$ , respectively, and the STELIB stellar library (Le Borgne et al., 2003), supplemented blueward of 3200 Å and redward 9500 Å by BaSel 3.1 spectra (Westera et al., 2002). The UV coverage of these SSPs permits computation of the  $Ly_c$  photon output as a function of time (see Sect. B.3). In computing the nebular continuum contribution we assumed an electron temperature and density of  $T_e = 10^4$  K and  $n_e = 100$  cm $^{-3}$ , respectively. Balmer emission-lines were computed from the total UV ionizing flux using the effective recombination coefficient  $\alpha_{H\alpha}^{eff}$  and by assuming ionization-bound nebulae (case B recombination).

Figure A.1 (upper panel) shows the time evolution of the  $EW(H\alpha)$  for four parametrizations of the SFH: instantaneous burst, continuous SF and exponentially decreasing SF with an e-folding time of 1 Gyr and 3 Gyr (black, yellow, green and violet curve, respectively). The  $EW(H\alpha)$  expected from the pAGB ( $\geq 10^8$  yr) stellar component for the combination of these models is displayed as the cyan shaded area. The lower-left panel of the figure shows the time evolution of the  $Ly_c$  photon rate per per solar mass, with the fractional contribution of the pAGB component displayed separately in the lower-right diagram.

From Fig. A.1 it is apparent the well-known fact (see, e.g., Leitherer et al., 1999; Weilbacher & Fritze-von Alvensleben, 2001; Kotulla et al., 2009) that the  $Ly_c$  photon rate and the corresponding Balmer-line EWs show a strong evolution with time, depending on the assumed SFH. For an instantaneous burst model, the  $Ly_c$  photon rate ( $s^{-1}$ ), following a steep decrease by  $\sim 5$  dex over the first  $10^8$  yr of galaxy evolution, levels off to a nearly constant level, scaling only with the available stellar mass as  $\approx 10^{41} M_\star/M_\odot$  (see also Cid Fernandes et al., 2011). This results in a nearly constant  $EW(H\alpha)$  in the narrow range between  $\sim 0.1$  Å and  $\sim 2.4$  Å ( $EW_\star^+$ ) over nearly the whole Hubble time. Note that an  $EW(H\alpha) \approx 0.1$  Å corresponds to a short ( $< 50$  Myr) period at the onset of the pAGB phase. As the curves within the shaded area of Fig. A.1 depict, the  $EW(H\alpha)$  predicted for the pure pAGB component for other SFHs also shows little evolution with time, with the notable trend, however, of remaining lower than that predicted for the instantaneous burst model. This is particularly true for the continuous SF model, which implies for the  $EW(H\alpha)$  owing to the pAGB component alone a by a




**Fig. A.1.** upper panel: Predicted  $EW(H\alpha)$  (Å) as a function of time for a solar-metallicity stellar population with a Salpeter IMF forming instantaneously (black), continuously at a constant SFR (yellow) and with an exponentially decreasing SFR with e-folding times  $t_e$  of 1 Gyr and 3 Gyr (green and red curve, respectively). The light-grey shaded area in the lower-right part of the diagram encompasses the range in  $EW(H\alpha)$  predicted from pAGB photoionization for ages  $\geq 0.1$  Gyr, and the narrow dark strip depicts the corresponding prediction ( $EW(H\alpha) \approx 1$  Å  $\equiv \langle EW_\star \rangle$ ) for the typical age of ETGs ( $\geq 6$  Gyr). The lower panel shows the time evolution of the  $Ly_c$  photon rate expected by the instantaneous and exponentially decreasing SFR models, and the mass fraction in % of stars older than 100 Myr (referred to as pAGB) is displayed separately in the lower-right diagram.

factor  $\sim 2$  lower value than the single-burst model. However, by considering the  $EW(H\alpha)$  expected to arise from the total (young+pAGB) stellar component (curves in the upper part of the same panel), it is obvious that continuous as well as exponentially decreasing ( $t_e=3$  Gyr) SF models can be discarded for ETGs, as they predict a large ( $\geq 6$  Å)  $EW(H\alpha)$  even for an age of  $\sim 10$  Gyr, whereas the observations (cf, e.g., P13 and Sect. 4) indicate values in the range of 1–2 Å. On the other hand, a model involving a shorter ( $t_e=1$  Gyr) phase of exponentially declining SFR (green curve) is compatible with the literature data, as it predicts an  $EW(H\alpha) \approx EW_\star^+$  after 6.3 Gyr of galactic evolution, just like the instantaneous SF model. For reasonable assumptions on the age of ETGs ( $\geq 6$  Gyr) these last two SFHs imply consistently a narrow span between  $\sim 1$  Å ( $\langle EW_\star \rangle$ ) and  $EW_\star^+$  for the  $EW(H\alpha)$  of photoionized gas. Allowing for a generous low limit of  $EW_\star^+=0.5$  Å (corresponding to a minimum  $\tau=2$  defined in P13 as a conservative low limit for the presence of  $Ly_c$  escape) we consider an observed  $EW(H\alpha)$  in the range between 0.5 Å and 2.4 Å to be consistent with the pAGB photoionization hypothesis.

## Appendix B: Analysis methodology

Figure B.1 provides a schematic overview of the three main modules of PORTO3D (v.2), which are briefly described next.



**Fig. B.1.** Schematic outline of the three main modules of PORTO3D (v.2).

### B.1. Module m.1: Computation of stellar fits and post-processing of the results

The module m.1, which is invoked at the first stage, extracts spectra from an IFS cube spaxel-by-spaxel, along with auxiliary information provided by the CALIFA data reduction pipeline (error spectrum, bad pixel map), computes initial statistics, and performs a number of preparatory steps that are necessary for subsequent fitting with STARLIGHT. These include the computation of the signal-to-noise (S/N) in various spectral windows (with the option of rejecting spaxels that do not satisfy certain quality criteria), the flagging of spurious features and emission lines within pre-defined or automatically generated spectral masks, and the spectral resampling to the restframe and wavelength rebinning to 1 Å/pixel. The observed flux at each wavelength  $O_\lambda$  and its uncertainty  $e_\lambda$ , provided by the CALIFA v1.3c reduction pipeline output or auto-computed with a sliding boxcar filter<sup>4</sup>, are finally exported together with flagging tags into ascii files in the format required for subsequent fitting with STARLIGHT. This module also creates up to 16 execution scripts, facilitating automated parallel spectral fitting of the data with multi-core CPUs. For each sample galaxy, typically  $\sim 2700$  individual spectra with a  $S/N \gtrsim 30$  at the normalization wavelength (5150 Å) were extracted and modeled with STARLIGHT.

The goal of STARLIGHT is the decomposition of an observed spectrum into a linear combination of its constituent SSP spectra. A description of the code is given in Cid Fernandes et al.

<sup>4</sup> This approach has been adopted in our pilot study of the ETGs NGC 5966 and NGC 6762 in K12, as an error spectrum was not provided in earlier versions of the CALIFA data reduction pipeline.

2005 (C05), 2007; and subsequent publications of the SEAGal collaboration<sup>5</sup> and is supplemented by a detailed cookbook for its application.

The best-fitting set of SSPs, together with the derived intrinsic V band extinction and velocity dispersion of the stellar component are referred to in the following as population vector. This encapsulates all information of the best-fitting model spectrum  $M_\lambda$

$$M_\lambda = M_{\lambda_0} \left[ \sum_{j=1}^{N_\star} x_j b_{j,\lambda} r_\lambda \right] \otimes G(v_\star, \sigma_\star),$$

where  $b_{j,\lambda} \equiv L_\lambda^{SSP}(t_j, Z_j) / L_{\lambda_0}^{SSP}(t_j, Z_j)$  is the spectrum of the  $j^{\text{th}}$  SSP normalized at  $\lambda_0$ ,  $r_\lambda \equiv 10^{-0.4(A_\lambda - A_{\lambda_0})}$  is the reddening term,  $\mathbf{x}$  is the population vector,  $M_{\lambda_0}$  is the synthetic flux at the normalization wavelength,  $N_\star$  is the total number of SSPs,  $G(v_\star, \sigma_\star)$  is the line-of-sight stellar velocity distribution, modeled as a Gaussian centered at velocity  $v_\star$  and broadened by  $\sigma_\star$ .

The convergence of the solution is driven by the standard  $\chi^2$  minimization with respect to the observed spectrum  $O_\lambda$

$$\chi^2(x, M_{\lambda_0}, A_V, v_\star, \sigma_\star) = \sum_{\lambda=1}^{N_\lambda} [(O_\lambda - M_\lambda) w_\lambda]^2,$$

whereby spectral weights  $w_\lambda$  are given by the inverse of the error spectrum  $e_\lambda$ .

STARLIGHT fits were computed in the spectral region between 4000 Å and 6800 Å. This was mainly because the S/N of our sample ETGs bluewards of 4000 Å is generally too low

<sup>5</sup> STARLIGHT & SEAGal: <http://www.starlight.ufsc.br/>



for a reliable stellar fit and the subsequent determination of the flux in the [OII] $\lambda\lambda 3726, 3729$  line doublet. The red spectral region (6900–7300 Å) was also disregarded because of its generally low S/N and spectral artifacts due to vignetting in the outer part of the PPAK FOV (see [Sánchez et al., 2012](#)).


For each spaxel STARLIGHT fits employing between 7 and 12 Markov chains were computed twice, using two different Simple Stellar Population (SSP) libraries, each comprising 34 ages (between 5 Myr and 13 Gyr) and three metallicities (0.008, 0.019, 0.03), i.e. 102 SSPs in total. The first set of models uses SSPs from [Bruzual & Charlot \(2003, hereafter BC03\)](#). As these cover the UV spectral range, they allow to infer the  $Ly_c$  photon output from each best-fitting population vector (see Sect. A for further details) which is in turn used for the computation of the  $\tau$  ratio (module m.3). For the second set of STARLIGHT fits we used SSPs from MILES ([Sánchez-Blázquez et al., 2006; Vazdekis et al., 2010; Falcón-Barroso et al., 2011](#)) which cover the optical spectral range (3540 – 7410 Å) only.

The module m.1 also permits spaxel binning prior to fitting according to various prescriptions (e.g., averaging to a constant S/N along azimuthal segments within isophotal annuli, see discussion in Sect. B.4 and Fig. B.5). Whereas a number of parallel runs on binned data cubes were conducted to check the robustness of the detection of faint nebular emission in the extranuclear component of our sample ETGs, we resorted to the methodology in K12 and P13 of carrying the spectral analysis in a single-spaxel (sisp) mode, and subsequently combining the results within irregular annuli (Sect. B.4).

Figure B.2 shows a high-quality (mean percentual deviation  $ADEV \approx 1.4$ ; cf STARLIGHT manual and [Cid Fernandes et al., 2005](#)) fit to a high-S/N spectrum from an off-center spaxel in the ETG NGC 5966. The observed spectrum  $O_\lambda$ , normalized at 5150 Å, and the fit ( $M_\lambda$ ) to it are shown in orange and blue color, respectively, in panel a. The lower-left panel b displays the fit residuals, with shaded regions depicting spectral regions flagged prior to the fit, or subsequently rejected by STARLIGHT. Panels c&d show, respectively, the luminosity ( $x_j$ ) and mass ( $\mu_j$ ) contribution of the individual SSPs ( $1 \dots j$ ) composing the best-fitting population vector. The thin grey vertical lines in the upper-right panel mark the 34 ages available in the SSP library for the three stellar metallicities (each coded by a different color), and the shaded area in the lower panel shows a smoothed version of the  $\mu_j$  distribution. The latter represents an illustrative approximation to the true SFH of stellar populations within the spaxel under study. It can be seen that the SFH involves an extended star formation episode occurring until  $\sim 7$  Gyr ago with a recent SF peak at  $\sim 10$  Myr. This fit therefore indicates that the stellar mass  $M_\star$  is dominated by an evolved pAGB stellar population with a mass-weighted age of 8.2 Gyr. It also implies little intrinsic extinction ( $\lesssim 0.2$  V mag) and no significant star-forming activity over the past  $\sim 1$  Gyr.

The module m.1, besides storing the relevant output from each fit ( $\sigma_\star$ ,  $v_\star$ , ..., among others), computes various quantities of interest, such as the time at which 50% of the present-day and ever-formed  $M_\star$  was in place, the luminosity contribution of stars formed in the past 5 Gyr ( $M_{\star,5\text{Gyr}}$ ), and the luminosity- and mass-weighted mean stellar age ( $\langle t_\star \rangle_L$  and  $\langle t_\star \rangle_M$ , respectively) as

$$\langle \log t_\star \rangle_L = \sum_{j=1}^{N_\star} x_j \log t_j, \quad (\text{B.1})$$



**Fig. B.3.** Luminosity-weighted and mass-weighted stellar age (upper and lower map, respectively) in Gyr (right-hand vertical bar) determined from STARLIGHT fits to individual spaxels. A trend for decreasing age with radius is apparent, in qualitative agreement with the results by [González Delgado et al. \(2014a\)](#). The  $\langle t_\star \rangle_M$  map indicates a mass-weighted stellar age of  $\geq 7$  Gyr all over the optical extent of the ETG.

and

$$\langle \log t_\star \rangle_M = \sum_{j=1}^{N_\star} \mu_j \log t_j. \quad (\text{B.2})$$


The dispersion of these quantities is also computed in the standard manner and exported into a data cube with dimension  $x \times y \times z$ , where  $x \times y$  is the dimension of the CALIFA input IFS cubes in arcsec, and  $z$  holds various quantities (age, metallicity, kinematics, emission-line fluxes, etc.) for each spaxel.

The  $\langle t_\star \rangle_M$  and  $M_{\star,5\text{Gyr}}$  maps provided by PORTO3D (see Fig. B.3 for an example) were only used for checking the soundness of the STARLIGHT fits and checking the pAGB nature of the stellar component in the studied ETGs.

## B.2. Module m.2: Computation of emission-line maps

The extraction of emission line fluxes in ETGs is in itself a challenging task due to their faintness, as already pointed out in several previous studies (e.g., [Sarzi et al., 2006; Annibali et al., 2010, K2, P13](#)). Indeed, faint ( $EW \sim 1$  Å) emission lines, buried within broader stellar absorption features, can only be detected and quantitatively studied upon precise subtraction of a high-S/N stellar continuum. Even in that case, a careful assessment of their uncertainties in individual spaxels is fundamental.

Previous longslit and IFS studies have generally investigated the diffuse *wim* component of ETGs down to EW levels of  $\geq 2$  Å,



**Fig. B.2.** a: Example of a high-quality fit with STARLIGHT to a V500 spectrum (blue and orange color, respectively) extracted from an off-nucleus spaxel of the ETG NGC 5966. From the lower-left panel b it is apparent that the fit residuals are small (on average,  $\lesssim 5\%$  of the stellar continuum level) yet systematic (see also Fig. B.4 and discussion in Sect. B.2). Shaded areas depict wavelength intervals that were flagged prior to spectral fitting. Panels c and d display, respectively, the luminosity ( $x_j$ ) and mass ( $\mu_j$ ) contribution of individual SSPs ( $1 \dots j$ ) in the best-fitting population vector. The thin grey vertical lines in the upper-right panel depict the 34 ages available in the SSP library for three stellar metallicities (color coded), and the shaded area in the lower panel shows a smoothed version of the  $\mu_j$ , which is meant as an illustration of the star formation history. The fit implies a dominant old (pAGB) stellar population suffering little intrinsic extinction ( $A_V \lesssim 0.2$  mag).


i.e. of the order of the EWs of Balmer stellar absorption profiles, leading to the conclusion that 2/3 of ETGs show faint nuclear and extranuclear nebular emission. A key contribution to advancing our understanding of the *wim* component in ETGs has been made by the SAURON project, which has first disclosed the diversity of emission-line morphologies and kinematics in these systems. Quite importantly, Sarzi et al. (2006) reported the detection of very faint ( $\text{EW} \sim 0.1 \text{ \AA}$ )  $\text{H}\beta$  emission in some ETGs, showing that even systems that were previously thought to be entirely devoid of ionized gas actually contain a low-surface brightness substrate of *wim*.

It is important to bear in mind, on the other hand, that emission-line flux determinations in ETGs are sensitively dependent on the quality of stellar fits, which in many cases show subtle yet systematic residuals. An example is given in Fig. B.4 where we show a typical, medium-quality fit to a *sisp* high-S/N spectrum. From the upper panel it can be seen that fitting residuals are minor (on average,  $\leq 5\%$  of the normalized continuum), yet systematic with both narrow ( $\sim 40 \text{ \AA}$ ) and broader ( $\sim 200 \text{ \AA}$ ) spectral intervals imperfectly fitted. These residuals were discussed in (e.g., Sarzi et al., 2006; Walcher et al., 2006, among others) and ascribed to a combination of various causes (e.g., imperfections in SSPs and/or fitting codes,  $\alpha$  element enhancement, contamination from telluric line emission). In the framework of CALIFA this issue was further analyzed in Husemann et al. (2013) through median-averaging of several thousands of individual spectra, showing indeed a systematic behavior in the residuals, in agreement with a parallel study by our team. The latter has shown that broad ripples in fitting residuals are partly to inaccuracies in the spectral sensitivity function adopted in the CALIFA v1.3c reduction pipeline. Indeed, a reprocessing with PORTO3D of part of the IFS data after reduction with the up-

graded version v1.4 shows that these broad residuals are much reduced or absent, although smaller-scale residuals generally persist. This issue is currently under scrutiny within the CALIFA collaboration.

On the other hand, as pointed out in Husemann et al. (2013), since these residuals are of the order or lower than the continuum noise level, their impact on emission line determinations is in practice negligible. Whereas we concur with this statement, based on the analysis of a few emission-line galaxies in our comparison sample (see Appendix D), we find that small-scale fitting residuals can significantly impact flux determinations for faint ( $\text{EW} \sim 1 \text{ \AA}$ ) emission lines when the standard technique of approximating the local rest-continuum by a first-order polynomial is used. In this case, Gaussian line fitting/deblending may fail to converge, or could lead to large errors owing to small-scale variations of the local background beneath and adjacent to emission lines. For this reason, these systematic fitting residuals could effectively place an EW limit of  $> 2 \text{ \AA}$  on studies of the *wim*, even when high-S/N spectra are available.

To alleviate this problem, PORTO3D performs for each *sisp* a rectification of the rest-continuum (i.e.  $O_\lambda - M_\lambda$ ), which is based on an 1D adaptation of the flux-conserving unsharp masking technique by Papaderos et al. (1998). This approach permits determination of a smooth version for the rest-continuum which is then subtracted from the input data array, yielding a net emission-line spectrum with a mean value of zero unless emission lines are present (cf lower panel of Fig. B.4). Other than the more conventional approach of re-fitting the residual continuum with a high-order polynomial, this technique captures and corrects both for large and small-scale fitting residuals, while leaving noise statistics unaffected.



**Fig. B.4.** upper panel: Subtraction of the best-fitting stellar SED computed with STARLIGHT with two different SSP libraries (BC03 and MILES; blue and magenta color, respectively) from an observed *sis*p spectrum extracted from an off-nucleus region of a CALIFA ETG. It can be seen that the fit residuals are comparatively small, of the order of  $1\text{--}2\sigma$  of the continuum in the blue spectral range, yet systematic. lower panel: The same spectrum after rectification of the rest-continuum.

The determination of emission-line fluxes from rectified net spectra is based on the combined application of two methods. The first one (e1) employs standard Gaussian line fitting/deblending with the Levenberg-Marquardt non-linear fitting algorithm and the second one (e2) a simple summation of the flux within pre-defined spectral windows centered on the laboratory wavelength of up to 50 emission lines, after correction for local motions. In the first case, the previous rectification of the residual continuum to a zero level is a key advantage, facilitating fast convergence and enhancing the robustness of emission line measurements, thereby permitting to extend the analysis down to  $\text{EWs} \sim 0.5 \text{ \AA}$  for a typical  $\text{S/N} \gtrsim 50$  in the continuum. This is largely because the continuum rectification effectively eliminates two free parameters (level and slope of the local continuum) in fitting/deblending. In cases where the deblending of the  $\text{H}\alpha$  line from the [NII] lines has failed to converge into a sensible solution, e.g., the [NII] $\lambda 6584/\text{H}\alpha$  ratio deviates from  $\sim 3$  beyond certain tolerance bounds, e1 switches to a coarse determination of the  $\text{H}\alpha$  flux by fitting the [NII] $\lambda 6584$  line only, and assuming a [NII] $\lambda 6584$ /[NII] $\lambda 6548 = 3$ . Prior to applying meth. e2, PORTO3D uses the gas kinematics pattern inferred from e1 to slightly adjust the center of the spectral windows  $[\lambda_{\min}:\lambda_{\max}]$  within which the flux summation is performed.

PORTO3D (v.2) estimates uncertainties in line fluxes by consideration in quadrature of four contributions to the error budget. The first one - taken into account only for e1 measurements - reflects formal errors in the fit, as obtained from the covariance matrix and is in most cases negligible. The second term considers photon statistics following a widely used formula (e.g., González-Delgado et al., 1995; Castellanos et al., 2002, see also K12)

$$\sigma = \sigma_{\text{cont}} N^{1/2} \left( 1 + \frac{\text{EW}}{N \Delta \lambda} \right)^{1/2} \quad (\text{B.3})$$

where  $\sigma_{\text{cont}}$  is the uncertainty in the determination of the local continuum (defined by its noise level),  $N$  the width of the region used to measure the line in pixels,  $\Delta \lambda$  the spectral dispersion in  $\text{\AA pix}^{-1}$ , and EW the equivalent width of the line in  $\text{\AA}$ .

Emission-line EWs were computed from the ratio between the line flux (measured on the net emission line spectra) and the rectified model SED ( $M_\lambda$ ). The third term, which is included in e2 measurements only, uses the noise distribution array in connection with an empirical scaling factor to gauge how the line flux registered within the spectral window considered could be affected by positive and negative photon noise peaks at the  $1\sigma$  level. Note that none of the above terms captures uncertainties related to the quality of the stellar fitting. An analytical estimate for the latter is obviously a non-trivial task, given that it has to also include a realistic description of how errors in the best-fitting stellar velocity dispersion and stellar metallicity may affect fluxes of emission lines being embedded within broader stellar absorption features. As a remedy to this problem, PORTO3D includes for e2 measurements an error contribution that is linked to the local deviation  $\delta_{\text{fit}}$  of the fit (i.e. the value of the rectification array within the respective spectral window  $\delta_\lambda$ ) as  $0.68 \cdot \delta_\lambda \cdot |\delta_{\text{fit}}|$ . This term, which typically dominates for faint emission-line fluxes, admittedly simplistic, is meant as a minimum attempt to quantify the cumulative impact of systematic errors in spectral fits on the recovered emission-line fluxes. Related to this issue is likely also the quality of the SSP models adopted. As we pointed out in K12, the fluxes of faint ( $\text{EW}(\text{H}\alpha) \lesssim 1 \text{ \AA}$ ) emission lines in ETGs may change to up to 10% depending on whether BC03 or MILES SSPs are used. For this reason, we have taken in this and our previous studies the computationally expensive approach of performing the full IFS data processing (spectral fitting and determination of emission lines) twice, both with the BC03 and MILES SSP library, and to subsequently combine the results. This was done with a decision-tree routine which prioritizes the up to four determinations for each spaxel (e1 and e2 for two SSP libraries) according to their soundness and estimated uncertainties. This procedure uses an initial sanity check of several quantities (errors, FWHM of emission lines, and in the case of the  $\text{H}\alpha$ + [NII] blend, the difference between their central wavelengths) and, in a second pass, the [NII] 6584/6548 ratios. Preference was generally given to e1 determinations, which, whenever available for



both BC- and MILES-based fits were error-weighted averaged. Whenever only one determination has passed the quality check, it was adopted for the final map, and the same decision procedure was followed for  $\text{H}\alpha$  measurements which were typically used in low-S/N, low-EW( $\text{H}\alpha$ ) spectra.

### B.3. Module m.3: Computation of the $\tau$ ratio

The  $\tau$  ratio, defined in K12 as the ratio of the Balmer-line luminosity expected to arise from pure pAGB photoionization to the observed one, offers a simple yet efficient means for assessing whether or not hot pAGB stars are capable of powering the extended *wim* in ETGs. After our pilot analysis of the radial distribution of  $\tau$  in two ETGs (K12), we extended this analysis in P13 to the present sample of 32 ETGs to test the pAGB photoionization hypothesis.

PORTO3D includes a routine that determines the total  $L_{\text{yc}}$  photon output for each STARLIGHT fit by integrating shortwards of the  $L_{\text{yc}}$  edge the UV continuum the SEDs of the individual SSPs (BC) composing the best-fitting population vector. In this procedure, only SSPs for stellar populations older than  $10^8$  yr are considered in order to compute the total  $L_{\text{yc}}$  output of the pAGB component only. The Balmer-line luminosities expected from it are computed assuming case B recombination and standard conditions for the gas ( $n_e = 100 \text{ cm}^{-3}$ ,  $T_e = 10^4 \text{ K}$ ). In ETGs, the typically large uncertainties in emission line fluxes and their ratios, generally prevent a reliable determination of the intrinsic extinction in the *wim*. Therefore, the optional inclusion of intrinsic extinction in PORTO3D m.3 has to rely on the stellar  $V$  band extinction, as inferred from STARLIGHT fits, which was generally found to be low ( $<0.1 \text{ mag}$ ).

As discussed in our previous studies (see also Sect. 1, a  $\tau \simeq 1$  corresponds to an equilibrium situation where the  $L_{\text{yc}}$  photon rate from pAGB stars precisely balances the observed  $\text{H}\alpha$  luminosity. A  $\tau < 1$ , on the other hand, indicates an additional contribution by an extra source, such as AGN, shocks or OB stars. Finally, as we discussed in P13, a  $\tau$  ratio exceeding unity corresponds to the situation of leakage of a fraction  $1-\tau^{-1}$  of the  $L_{\text{yc}}$  output from the pAGB component. However, given the inherent uncertainties in the currently available model predictions for the UV output from pAGB stars, we conservatively assumed in P13 and in this study a  $\tau \geq 2$  as robust evidence for  $L_{\text{yc}}$  escape.

### B.4. Computation of radial profiles

In Sect. 4 we show the radial distribution of various quantities of interest, such as the intensity and EW of the  $\text{H}\alpha$  emission line, and the  $[\text{O III}]_{5007}/\text{H}\beta$ ,  $[\text{N II}]_{6584}/\text{H}\alpha$  and  $\tau$  ratios. Following our methodology in P13, we include in all diagrams two complementary data sets:

i) single-spaxel (*sisp*) determinations from fits with an ADEV  $|O_\lambda - M_\lambda| / O_\lambda \leq 2.6$  (cf K12 and the STARLIGHT manual). This higher-S/N subsample has the lowest uncertainties in fitting parameters and emission-line fluxes at the price of being spatially restricted to typically the central, higher-surface brightness part ( $\mu \lesssim 23 \text{ g mag}/\square''$ ) of the studied galaxies.

ii) The average of all *sisp* determinations within isophotal annuli (*isan*) adapted to the morphology of the (line-free) continuum between 6390 Å and 6490 Å. These data, which are to be considered in a statistical sense, go  $\geq 2 \text{ mag}$  fainter, allowing the study of the *wim* in the ETG periphery.


The isophotal annuli method (Papaderos et al., 2002, referred to as meth. iv; subsequently as *Lazy* by Noeske et al. 2003,2006)



**Fig. B.5.** Adaptation of the irregular annuli surface photometry technique by Papaderos et al. (2002) for the purpose of azimuthal binning of spaxels to a constant S/N within radial zones. The color coding is meant to illustrate the segmentation within isophotal annuli.

was developed with the goal of the derivation of radial surface brightness and nebular line intensity profiles in irregular galaxies with a not well constrained optical center, for which the conventional approach of fitting ellipses to isophotes of 2D axis-symmetric luminosity components is of limited applicability. Its main concept consists in the determination of photon statistics within equidistant logarithmic slices of a galaxy image (or of a reference image, such as, e.g., a stacked image of all available passbands, or in the context of this study the line-free 6390–6490 Å pseudo-continuum image) with each of these irregular slices faithfully reproducing the morphology of a galaxy within each intensity interval  $[I: I+\delta I]$  and corresponding to a well-defined photometric radius  $R^*$ . In the context of IFS data, this technique was first used in K12 (see their Appendix for further details) and more recently by Sánchez et al. (2014b) for the derivation of gas-phase metallicity gradients in CALIFA late-type galaxies.


Figure B.5 illustrates the adaptation of the isophotal annuli technique such as to permit azimuthal binning of spaxels to a constant S/N within radial zones. This add-on routine to PORTO3D was used in P13 and in this study in order to double-check the presence of extended nebular emission in ETGs. These tests were supplemented through the modeling of the integrated spectrum within each annulus, extracted by the same routine by requesting an unrealistically high S/N. Figure B.6 shows examples of the residual emission after subtraction of the stellar fit with BC SSPs and continuum rectification within successively large annuli for three ETGs from our sample, illustrating the extreme faintness of nebular emission in type ii ETGs (NGC 6338 and NGC 7550; panels b&c) as compared to type i ETGs (NGC 1167; panel a).



**Fig. B.6.** Integrated spectra of the type i ETG NGC 1167 (a) and the type ii ETGs NGC 6338 and NGC 7550 (panels b and c, respectively) within successive isophotal annuli (isan), after subtraction of the stellar emission and rectification of the continuum. Each spectrum is labeled with the photometric radius (") of the annulus within which it was sampled and vertically shifted by an arbitrary amount, with the overlaid dotted horizontal lines depicting the zero level. The vertical bar in each panel corresponds to a flux level interval of  $10^{-16} \text{ erg s}^{-1} \text{ cm}^{-2}$ . The extreme faintness of extended nebular emission in type ii ETGs, as compared to the type i ETG NGC 1167 is apparent.

## Appendix C: 2D emission-line patterns and kinematics for 32 galaxies

This section illustrates the diversity of ETG galaxies within our sample with respect to morphology, kinematics and physical properties of their warm interstellar medium. The galaxies are ordered according to Table 1. Note that the maps and radial profiles (a–n) follow the same definitions and colors as in Sect. 4.



## UGC 5771

Morphology NED: S0/a

ETG type i

$\langle \tau(\text{H}\alpha)_e \rangle = 0.69$


$D = 106.4 \text{ Mpc} \mid M_r = -21.9 \text{ mag}$

$L(\text{H}\alpha) = 34.0 \times 10^{39} \text{ erg s}^{-1}$

$\log M_\star = 11.398 [M_\odot]$

**Fig. C.1.** UGC 5771: 2D maps (panels a-f and n), radial intensity and EW of  $\text{H}\alpha$  (panels g and h), BPT diagrams (panels i and j), radial distribution of diagnostic line ratios (k and l), and the  $\tau$  ratio (panel m). In all 2D maps north is up and east to the left. The horizontal bar corresponds to  $20''$ .





## NGC 4003

Morphology NED: SB0

ETG type i


$\langle \tau(\text{H}\alpha) \rangle = 0.57$

$D = 96.6 \text{ Mpc} \mid M_r = -21.71 \text{ mag}$

$L(\text{H}\alpha) = 50.8 \times 10^{39} \text{ erg s}^{-1}$

$\log M_\star = 11.218 [M_\odot]$

**Fig. C.2.** NGC 4003: 2D maps (panels a-f and n), radial intensity and EW of  $\text{H}\alpha$  (panels g and h), BPT diagrams (panels i and j), radial distribution of diagnostic line ratios (k and l), and the  $\tau$  ratio (panel m). In all 2D maps north is up and east to the left. The horizontal bar corresponds to  $20''$ .



## UGC 8234

Morphology NED: SO/a

ETG type i


$\langle \tau(\text{H}\alpha)_e \rangle = 1.09$

$D = 116.1 \text{ Mpc} \mid M_r = -22.5 \text{ mag}$

$L(\text{H}\alpha) = 15.9 \times 10^{39} \text{ erg s}^{-1}$


$\log M_\star = 11.131 [M_\odot]$

**Fig. C.3.** UGC 8234: 2D maps (panels a-f and n), radial intensity and EW of  $\text{H}\alpha$  (panels g and h), BPT diagrams (panels i and j), radial distribution of diagnostic line ratios (k and l), and the  $\tau$  ratio (panel m). In all 2D maps north is up and east to the left. The horizontal bar corresponds to  $20''$ .




**Fig. C.4.** NGC 5966: 2D maps (panels a-f and n), radial intensity and EW of  $\text{H}\alpha$  (panels g and h), BPT diagrams (panels i and j), radial distribution of diagnostic line ratios (k and l), and the  $\tau$  ratio (panel m). In all 2D maps north is up and east to the left. The horizontal bar corresponds to  $20''$ .






**Fig. C.5.** UGC 10205: 2D maps (panels a-f and n), radial intensity and EW of  $\text{H}\alpha$  (panels g and h), BPT diagrams (panels i and j), radial distribution of diagnostic line ratios (k and l), and the  $\tau$  ratio (panel m). In all 2D maps north is up and east to the left. The horizontal bar corresponds to  $20''$ .

The warm ionized gas in CALIFA early-type galaxies





## NGC 6146

Morphology NED: E?

ETG type i

$\langle \tau(H\alpha)_e \rangle = 1.86$

$D = 127.4 \text{ Mpc}$  |  $M_r = -23.04 \text{ mag}$


$L(H\alpha) = 34.9 \times 10^{39} \text{ erg s}^{-1}$

$\log M_\star = 11.720 [M_\odot]$


**Fig. C.7.** NGC 6146: 2D maps (panels a-f and n), radial intensity and EW of H $\alpha$  (panels g and h), BPT diagrams (panels i and j), radial distribution of diagnostic line ratios (k and l), and the  $\tau$  ratio (panel m). In all 2D maps north is up and east to the left. The horizontal bar corresponds to  $20''$ .



The warm ionized gas in CALIFA early-type galaxies



**Fig. C.8.** UGC 10695: 2D maps (panels a-f and n), radial intensity and EW of  $H\alpha$  (panels g and h), BPT diagrams (panels i and j), radial distribution of diagnostic line ratios (k and l), and the  $\tau$  ratio (panel m). In all 2D maps north is up and east to the left. The horizontal bar corresponds to  $20''$ .



## UGC 10905

Morphology NED: S0/a

ETG type i


$\langle \tau(\text{H}\alpha) \rangle = 0.98$

$D = 114.1 \text{ Mpc}$  |  $M_r = -22.26 \text{ mag}$


$L(\text{H}\alpha) = 33.8 \times 10^{39} \text{ erg s}^{-1}$

$\log M_{\star} = 11.487 [M_{\odot}]$

**Fig. C.9.** UGC 10905: 2D maps (panels a-f and n), radial intensity and EW of  $\text{H}\alpha$  (panels g and h), BPT diagrams (panels i and j), radial distribution of diagnostic line ratios (k and l), and the  $\tau$  ratio (panel m). In all 2D maps north is up and east to the left. The horizontal bar corresponds to  $20''$ .




**Fig. C.10.** NGC 6762: 2D maps (panels a-f and n), radial intensity and EW of  $H\alpha$  (panels g and h), BPT diagrams (panels i and j), radial distribution of diagnostic line ratios (k and l), and the  $\tau$  ratio (panel m). In all 2D maps north is up and east to the left. The horizontal bar corresponds to  $20''$ .



**Fig. C.11.** NGC 7025: 2D maps (panels a-f and n), radial intensity and EW of H $\alpha$  (panels g and h), BPT diagrams (panels i and j), radial distribution of diagnostic line ratios (k and l), and the  $\tau$  ratio (panel m). In all 2D maps north is up and east to the left. The horizontal bar corresponds to  $20''$ .





## NGC 1349

Morphology NED: S0

ETG type i+


$\langle \tau(\text{H}\alpha) \rangle = 0.89$

$D = 87.7 \text{ Mpc} \mid M_r = -21.62 \text{ mag}$

$L(\text{H}\alpha) = 42.9 \times 10^{39} \text{ erg s}^{-1}$

$\log M_{\star} = 11.183 [M_{\odot}]$

**Fig. C.12.** NGC 1349: 2D maps (panels a-f and n), radial intensity and EW of H $\alpha$  (panels g and h), BPT diagrams (panels i and j), radial distribution of diagnostic line ratios (k and l), and the  $\tau$  ratio (panel m). In all 2D maps north is up and east to the left. The horizontal bar corresponds to  $20''$ .



## NGC 3106

Morphology NED: S0

ETG type i+


$\langle \tau(\text{H}\alpha)_e \rangle = 0.60$

$D = 90.1 \text{ Mpc} \mid M_r = -22.08 \text{ mag}$


$L(\text{H}\alpha) = 63.0 \times 10^{39} \text{ erg s}^{-1}$

$\log \mathcal{M}_\star = 11.314 [M_\odot]$

**Fig. C.13.** NGC 3106: 2D maps (panels a-f and n), radial intensity and EW of H $\alpha$  (panels g and h), BPT diagrams (panels i and j), radial distribution of diagnostic line ratios (k and l), and the  $\tau$  ratio (panel m). In all 2D maps north is up and east to the left. The horizontal bar corresponds to  $20''$ .



**Fig. C.14.** UGC 0029: 2D maps (panels a-f and n), radial intensity and EW of  $H\alpha$  (panels g and h), BPT diagrams (panels i and j), radial distribution of diagnostic line ratios (k and l), and the  $\tau$  ratio (panel m). In all 2D maps north is up and east to the left. The horizontal bar corresponds to  $20''$ .



## NGC 2918

Morphology NED: E

ETG type ii

$\langle \tau(\text{H}\alpha) \rangle = 4.87$

$D = 98.1 \text{ Mpc} \mid M_r = -22.40 \text{ mag}$


$L(\text{H}\alpha) = 6.9 \times 10^{39} \text{ erg s}^{-1}$

$\log M_\star = 10.340 [M_\odot]$


**Fig. C.15.** NGC 2918: 2D maps (panels a-f and n), radial intensity and EW of  $\text{H}\alpha$  (panels g and h), BPT diagrams (panels i and j), radial distribution of diagnostic line ratios (k and l), and the  $\tau$  ratio (panel m). In all 2D maps north is up and east to the left. The horizontal bar corresponds to  $20''$ .



The warm ionized gas in CALIFA early-type galaxies




**Fig. C.16.** NGC 3300: 2D maps (panels a-f and n), radial intensity and EW of H $\alpha$  (panels g and h), BPT diagrams (panels i and j), radial distribution of diagnostic line ratios (k and l), and the  $\tau$  ratio (panel m). In all 2D maps north is up and east to the left. The horizontal bar corresponds to  $20''$ .




**Fig. C.17.** NGC 3615: 2D maps (panels a-f and n), radial intensity and EW of  $\text{H}\alpha$  (panels g and h), BPT diagrams (panels i and j), radial distribution of diagnostic line ratios (k and l), and the  $\tau$  ratio (panel m). In all 2D maps north is up and east to the left. The horizontal bar corresponds to  $20''$ .

The warm ionized gas in CALIFA early-type galaxies




**Fig. C.18.** NGC 4816: 2D maps (panels a-f and n), radial intensity and EW of H $\alpha$  (panels g and h), BPT diagrams (panels i and j), radial distribution of diagnostic line ratios (k and l), and the  $\tau$  ratio (panel m). In all 2D maps north is up and east to the left. The horizontal bar corresponds to  $20''$ .




**Fig. C.19.** NGC 6125: 2D maps (panels a-f and n), radial intensity and EW of  $\text{H}\alpha$  (panels g and h), BPT diagrams (panels i and j), radial distribution of diagnostic line ratios (k and l), and the  $\tau$  ratio (panel m). In all 2D maps north is up and east to the left. The horizontal bar corresponds to  $20''$ .



The warm ionized gas in CALIFA early-type galaxies





## NGC 6173

Morphology NED: E

ETG type ii

$\langle \tau(\text{H}\alpha) \rangle = 15.6$


$D = 126.9 \text{ Mpc} \mid M_r = -23.19 \text{ mag}$

$L(\text{H}\alpha) = 8.6 \times 10^{39} \text{ erg s}^{-1}$

$\log M_\star = 11.999 [M_\odot]$

**Fig. C.21.** NGC 6173: 2D maps (panels a-f and n), radial intensity and EW of  $\text{H}\alpha$  (panels g and h), BPT diagrams (panels i and j), radial distribution of diagnostic line ratios (k and l), and the  $\tau$  ratio (panel m). In all 2D maps north is up and east to the left. The horizontal bar corresponds to  $20''$ .

The warm ionized gas in CALIFA early-type galaxies



## UGC 10693

Morphology NED: E

ETG type ii


$\langle \tau(\text{H}\alpha)_e \rangle = 11.7$

$D = 120.5 \text{ Mpc} \mid M_r = -22.70 \text{ mag}$

$L(\text{H}\alpha) = 5.9 \times 10^{39} \text{ erg s}^{-1}$

$\log \mathcal{M}_* = 11.572 [\text{M}_\odot]$

**Fig. C.22.** UGC 10693: 2D maps (panels a-f and n), radial intensity and EW of H $\alpha$  (panels g and h), BPT diagrams (panels i and j), radial distribution of diagnostic line ratios (k and l), and the  $\tau$  ratio (panel m). In all 2D maps north is up and east to the left. The horizontal bar corresponds to  $20''$ .



## NGC 6338

Morphology NED: S0

ETG type ii

$\langle \tau(\text{H}\alpha) \rangle = 2.21$

$D = 117.5 \text{ Mpc}$  |  $M_r = -22.79 \text{ mag}$


$L(\text{H}\alpha) = 29.4 \times 10^{39} \text{ erg s}^{-1}$

$\log \mathcal{M}_\star = 11.707 [\text{M}_\odot]$


**Fig. C.23.** NGC 6338: 2D maps (panels a-f and n), radial intensity and EW of  $\text{H}\alpha$  (panels g and h), BPT diagrams (panels i and j), radial distribution of diagnostic line ratios (k and l), and the  $\tau$  ratio (panel m). In all 2D maps north is up and east to the left. The horizontal bar corresponds to  $20''$ .



The warm ionized gas in CALIFA early-type galaxies




**Fig. C.24.** NGC 6411: 2D maps (panels a-f and n), radial intensity and EW of  $\text{H}\alpha$  (panels g and h), BPT diagrams (panels i and j), radial distribution of diagnostic line ratios (k and l), and the  $\tau$  ratio (panel m). In all 2D maps north is up and east to the left. The horizontal bar corresponds to  $20''$ .



**Fig. C.25.** NGC 6427: 2D maps (panels a-f and n), radial intensity and EW of  $\text{H}\alpha$  (panels g and h), BPT diagrams (panels i and j), radial distribution of diagnostic line ratios (k and l), and the  $\tau$  ratio (panel m). In all 2D maps north is up and east to the left. The horizontal bar corresponds to  $20''$ .

The warm ionized gas in CALIFA early-type galaxies



## NGC 6515

Morphology NED: E

ETG type ii


$\langle \tau(\text{H}\alpha) \rangle = 2.46$

$D = 99.0 \text{ Mpc} \mid M_r = -22.11 \text{ mag}$

$L(\text{H}\alpha) = 10.2 \times 10^{39} \text{ erg s}^{-1}$

$\log M_\star = 11.316 [M_\odot]$

**Fig. C.26.** NGC 6515: 2D maps (panels a-f and n), radial intensity and EW of  $\text{H}\alpha$  (panels g and h), BPT diagrams (panels i and j), radial distribution of diagnostic line ratios (k and l), and the  $\tau$  ratio (panel m). In all 2D maps north is up and east to the left. The horizontal bar corresponds to  $20''$ .



## NGC 7194

Morphology NED: E

ETG type ii

$\langle \tau(\text{H}\alpha) \rangle = 12.9$

$D = 110.7 \text{ Mpc} \mid M_r = -22.35 \text{ mag}$


$L(\text{H}\alpha) = 4.4 \times 10^{39} \text{ erg s}^{-1}$

$\log \mathcal{M}_\star = 11.536 [M_\odot]$

**Fig. C.27.** NGC 7194: 2D maps (panels a-f and n), radial intensity and EW of  $\text{H}\alpha$  (panels g and h), BPT diagrams (panels i and j), radial distribution of diagnostic line ratios (k and l), and the  $\tau$  ratio (panel m). In all 2D maps north is up and east to the left. The horizontal bar corresponds to  $20''$ .



The warm ionized gas in CALIFA early-type galaxies



## NGC 7236

Morphology NED: SA0

ETG type ii


$\langle \tau(\text{H}\alpha) \rangle_e = 8.16$

$D = 108.2 \text{ Mpc} \mid M_r = -21.45 \text{ mag}$

$L(\text{H}\alpha) = 5.0 \times 10^{39} \text{ erg s}^{-1}$


$\log \mathcal{M}_* = 11.403 [M_\odot]$

**Fig. C.28.** NGC 7236: 2D maps (panels a-f and n), radial intensity and EW of  $\text{H}\alpha$  (panels g and h), BPT diagrams (panels i and j), radial distribution of diagnostic line ratios (k and l), and the  $\tau$  ratio (panel m). In all 2D maps north is up and east to the left. The horizontal bar corresponds to  $20''$ .




**Fig. C.29.** UGC 11958: 2D maps (panels a-f and n), radial intensity and EW of  $\text{H}\alpha$  (panels g and h), BPT diagrams (panels i and j), radial distribution of diagnostic line ratios (k and l), and the  $\tau$  ratio (panel m). In all 2D maps north is up and east to the left. The horizontal bar  $50''$  corresponds to  $20''$ .

The warm ionized gas in CALIFA early-type galaxies



**Fig. C.30.** NGC 7436B: 2D maps (panels a-f and n), radial intensity and EW of  $H\alpha$  (panels g and h), BPT diagrams (panels i and j), radial distribution of diagnostic line ratios (k and l), and the  $\tau$  ratio (panel m). In all 2D maps north is up and east to the left. The horizontal bar corresponds to  $20''$ .




**Fig. C.31.** NGC 7550: 2D maps (panels a-f and n), radial intensity and EW of  $\text{H}\alpha$  (panels g and h), BPT diagrams (panels i and j), radial distribution of diagnostic line ratios (k and l), and the  $\tau$  ratio (panel m). In all 2D maps north is up and east to the left. The horizontal bar  $\varnothing$  corresponds to  $20''$ .




## **Appendix D: Comparison with star-forming/composite galaxies**


This appendix is meant to illustrate the variation of the  $\text{EW}(\text{H}\alpha)$ ,  $\tau$ ; and diagnostic line ratios from early-type towards late-type galaxies. All but one of these systems are classifiable by BPT ratios in their central part and/or extranuclear component as *composite* or *SF-dominated*.



**Fig. D.1. IC 540:** 2D maps (panels a-f and n), radial intensity and EW of  $\text{H}\alpha$  (panels g and h), BPT diagrams (panels i and j), radial distribution of diagnostic line ratios (k and l), and the  $\tau$  ratio (panel m). In all 2D maps north is up and east to the left. The horizontal bar corresponds to  $20''$ .




**Fig. D.2.** UGC 8778: 2D maps (panels a–f and n), radial intensity and EW of  $H\alpha$  (panels g and h), BPT diagrams (panels i and j), radial distribution of diagnostic line ratios (k and l), and the  $\tau$  ratio (panel m). In all 2D maps north is up and east to the left. The horizontal bar corresponds to  $20''$ .




**Fig. D.3. IC 944:** 2D maps (panels a-f and n), radial intensity and EW of  $H\alpha$  (panels g and h), BPT diagrams (panels i and j), radial distribution of diagnostic line ratios (k and l), and the  $\tau$  ratio (panel m). In all 2D maps north is up and east to the left. The horizontal bar corresponds to  $20''$ .



The warm ionized gas in CALIFA early-type galaxies



**Fig. D.4.** IC 1683: 2D maps (panels a-f and n), radial intensity and EW of  $\text{H}\alpha$  (panels g and h), BPT diagrams (panels i and j), radial distribution of diagnostic line ratios (k and l), and the  $\tau$  ratio (panel m). In all 2D maps north is up and east to the left. The horizontal bar corresponds to  $20''$ .



## NGC 4470

### Comparison Sample

Morphology NED: Sa?

$\langle \tau(\text{H}\alpha)_e \rangle = 0.03$

$D = 38.4 \text{ Mpc} \mid M_r = -20.35 \text{ mag}$

$L(\text{H}\alpha) = 147 \times 10^{39} \text{ erg s}^{-1}$

$\log \mathcal{M}_\star = 10.108 [M_\odot]$

**Fig. D.5.** NGC 4470: 2D maps (panels a-f and n), radial intensity and EW of H $\alpha$  (panels g and h), BPT diagrams (panels i and j), radial distribution of diagnostic line ratios (k and l), and the  $\tau$  ratio (panel m). In all 2D maps north is up and east to the left. The horizontal bar corresponds to  $20''$ .

- 
- <sup>1</sup> Instituto de Astrofísica e Ciências do Espaço, Universidade do Porto, Centro de Astrofísica da Universidade do Porto, Rua das Estrelas, 4150-762 Porto, Portugal
  - <sup>2</sup> Instituto de Astrofísica de Andalucía (CSIC), Glorieta de la Astronomía s/n Aptdo. 3004, E18080-Granada, Spain
  - <sup>3</sup> Institut d'Astrophysique de Paris, UMR 7095, CNRS, Université Pierre et Marie Curie, 98 bis boulevard Arago, 75014 Paris, France
  - <sup>4</sup> Centro Astronómico Hispano Alemán de Calar Alto (CSIC-MPG), C/ Jesús Durbán Remón 2-2, E-4004 Almería, Spain
  - <sup>5</sup> University of Vienna, Türkenschanzstrasse 17, 1180 Vienna, Austria
  - <sup>6</sup> Estación Experimental de Zonas Áridas (CSIC), Ctra. de Sacramento s.n., La Cañada, Almería, Spain
  - <sup>7</sup> Sydney Institute for Astronomy, University of Sydney, NSW 2006, Australia
  - <sup>8</sup> Millennium Institute of Astrophysics, Chile
  - <sup>9</sup> Departamento de Astronomía, Universidad de Chile, Casilla 36-D, Santiago, Chile
  - <sup>10</sup> Astronomical Institute of the Ruhr-University Bochum
  - <sup>11</sup> RUB Research Department Plasmas with Complex Interactions
  - <sup>12</sup> Instituto Nacional de Astrofísica, Óptica y Electrónica, Luis E. Erro 1, 72840 Tonantzinla, Puebla, Mexico
  - <sup>13</sup> Departamento de Física, Universidade Federal de Santa Catarina, PO Box 476, 88040-900, Florianópolis, SC, Brazil
  - <sup>14</sup> Leibniz-Institut für Astrophysik Potsdam (AIP), An der Sternwarte 16, D-14482 Potsdam, Germany
  - <sup>15</sup> Instituto de Astrofísica de Canarias, Vía Láctea s/n, La Laguna, Tenerife, Spain
  - <sup>16</sup> Departamento de Astrofísica, Universidad de La Laguna, E-38205 La Laguna, Tenerife, Spain
  - <sup>17</sup> CIEMAT, Avda. Complutense 40, 28040 Madrid, Spain
  - <sup>18</sup> CEI Campus Moncloa, UCM-UPM, Departamento de Astrofísica y CC. de la Atmósfera, Facultad de CC. Físicas, Universidad Complutense de Madrid, Avda. Complutense s/n, 28040 Madrid, Spain
  - <sup>19</sup> Department of Physics, Institute for Astronomy, ETH Zürich, CH-8093 Zürich, Switzerland
  - <sup>20</sup> Australian Astronomical Observatory, PO Box 915, North Ryde, NSW 1670, Australia
  - <sup>21</sup> Department of Physics and Astronomy, Macquarie University, NSW 2109, Australia